arXiv:1207.0414v1 [nucl-th] 2 Jul 2012

An alternative well founded way to treat the center-of-mass correlations: proposition of a local center-of-mass correlations potential.

Jérémie Messud[1]

[1] *Université Bordeaux, CNRS/IN2P3, Centre d'Etudes Nucléaires de Bordeaux Gradignan, UMR5797, F-33175 Gradignan, France*

(Dated: July 3, 2012)

The recently developped "internal" Density Functional Theory provides an existence theorem for a local potential that contains the center-of-mass correlations effects. The knowledge of the corresponding energy functional would provide a much cheaper way than projection techniques to treat those correlations. The aim of this article is to construct such a functional. We propose a well founded method, suitable for Fermions as well as for Bosons, which does not necessit to include any free parameter.

I. INTRODUCTION.

One of the most obvious symmetry of isolated self-bound systems (as atomic nuclei, Helium droplets or molecular systems where nuclei are treated explicitely) is translational invariance. Translational invariance of self-bound systems Hamiltonians ensures Galilean invariance of the wave function, so that the center-of-mass (c.m.) properties can be separated from the "internal" ones (that are always deduced by experimentalists). As a consequence, one laboratory coordinate is redundant for the description of the internal properties, which gives rise to c.m. correlations.

A numerically manageable and succesfull way to describe self-bound systems is to use mean-field like calculations with effective interactions. The corresponding equations are often justified starting from the Hartree-Fock (HF) framework, which sacrifices by construction "Galilean invariance for the sake of the Pauli principle", to quote Ref. [1]. As a consequence, the c.m. correlations (or equivalently the redundant coordinate problem) are (is) not treated correctly. This introduces a spurious coupling between the internal properties and the c.m. motion in a HF framework, that affects the energy and other observables [2].

A way to overcome this problem in the stationary case is to perform projected HF, where projection before variation on c.m. momentum restores translational or Galilean invariance of the wave-function. Peierls and Yoccoz proposed a (single) projection method to restore translational invariance [3]. Later, Peierls and Thouless proposed a (double) projection method to restore the more fundamental Galilean invarianve [4]. To our knowledge, every numerical calculations that treat the c.m. correlations by projection before variation have been done within the Peirls and Yoccoz method [5–8], thus not restoring the full Galilean invariance. Moreover, the prices to pay are to abandon the independent-particle description and a large numerical cost [1, 4, 9, 10]. Indeed, projection techniques require "an order of magnitude more computing time than the underlying mean-field like calculations", to quote Ref. [7], which is discriminatory for the description of intermediary size systems. This led to the development of various approximate methods to treat the c.m. correlations, see Ref. [11] for an overview. For instance, a common method is to add a $-<\frac{\mathbf{P}^2}{2mN}>$ term in the energy functional (more details will be given in §II C). But the successes of those methods are not systematic and the approximations done not perfectly justified.

In the time-dependent case, the spurious c.m. motion problem remains [12, 13], but the situation is more tricky because then the projected HF method becomes unmanageable even for very small self-bound systems [12]. It thus remains an open question to develop a rigorous and numerically cheap scheme to treat the c.m. correlations, which would go beyond standard approximations and would remain usable in the time-dependent case.

So far, the search for such a scheme has not been studied extensively, maybe because it is sometimes thought that the c.m. correlations problem concerns only very small self-bound systems. But the c.m correlations can have a non negligible effect even for intermediary size systems. For instance, it has been shown that c.m. correlations are not negligible for all nuclei above ^{16}O [6, 7, 14]. This reinforces the necessity to develop a numerically manageable method to treat those correlations.

A rigorous alternative and *a priori* numerically much less costly way to take into account the c.m. correlations has been opened by the recently developped "internal" Density Functional Theory (DFT) and Kohn-Sham (KS) scheme [15–17]. Differing from standard DFT [18–21], it is formulated in the c.m frame of a self-bound system and proves that the c.m. correlations can be included in the energy functional and thus in a local KS potential [15–17]. In addition to the fact that it gives a much more fundamental justification than the HF framework to the use of mean-field like calculations with effective interactions for the description of self-bound systems, it shows that there would be no need for a c.m. projection if the ultimate functional was known. Internal DFT gives an existence theorem but not a constructive method. The aim of the present article is to propose such a method.

Typeset by REVTEX

The article is organized as follows. In section II is provided a quick recall of the internal DFT formalism and underlined the limitations of the commonly used methods to treat the c.m. correlations. In section III is developped a new general form for a local c.m. correlations potential, whithout including any free parameter. Finally, in section IV are given preliminary but convincing numerical results on various model systems.

II. INTERNAL DFT AND THE C.M. CORRELATIONS FUNCTIONAL.

A. Quick recall of the internal DFT formalism.

We start from a self-bound system composed by N identical particles (Fermions or Bosons) and follow the considerations of Ref. [15]. The coordinates of the particles in any chosen inertial frame of reference (such as the laboratory) are denoted $\{\mathbf{r}_i\}$. The c.m. coordinate of the system is denoted

$$\mathbf{R} = \frac{1}{N}\sum_{j=1}^{N} \mathbf{r}_j .$$

The system is described by the following translationally invariant N-body Hamiltonian:

$$H = \sum_{i=1}^{N} \frac{\mathbf{p}_i^2}{2m} + \sum_{\substack{i,j=1 \\ i>j}}^{N} u(\mathbf{r}_i - \mathbf{r}_j) + \sum_{i=1}^{N} v^{\text{int}}(\mathbf{r}_i - \mathbf{R}) \, , \tag{1}$$

composed by the usual kinetic energy term, a 2-body potential u which describes the Fermion-Fermion (or Boson-Boson) interaction (generalization of the following considerations to 3-body... interactions is straightforward) and an arbitrary translationally invariant potential v^{int}. This last potential is an "internal" potential, i.e. it is defined in the c.m. frame an acts only on the internal properties. Of course it is zero in the purely isolated self-bound case. Nevertheless, its form is suitable to model internal effects of fields used in experiments (polarization potentials...) [16].

We introduce the Jacobi coordinates ξ_α defined as $\xi_1 = \mathbf{r}_2 - \mathbf{r}_1$, $\xi_2 = \mathbf{r}_3 - \frac{\mathbf{r}_2+\mathbf{r}_1}{2}$, ..., $\xi_{N-1} = \frac{N}{N-1}(\mathbf{r}_N - \mathbf{R})$. This permits us to separate the Hamiltonian (1) into $H = H_{CM} + H_{int}$, where $H_{CM} = -(\hbar^2/2M)\Delta_{\mathbf{R}}$ (with $M = Nm$ being the total mass) is a 1-body Hamiltonian dercribing the c.m. motion and acting in the $\mathbf{R}$ space only, and H_{int} is a $(N-1)$ body-Hamiltonian describing the internal properties and acting in the $\{\xi_\alpha\}$ space only:

$$H_{int} = \sum_{\alpha=1}^{N-1} \frac{\tau_\alpha^2}{2\mu_\alpha} + U(\xi_1, \ldots, \xi_{N-1}) + V^{\text{int}}(\xi_1, \ldots, \xi_{N-1}) \, . \tag{2}$$

H_{int} contains the interaction u and the potential v^{int}, because they can be rewritten as functions of the $\{\xi_\alpha\}$ only [denoted respectively $U(\xi_1, \ldots, \xi_{N-1})$ and $V^{\text{int}}(\xi_1, \ldots, \xi_{N-1})$], and the internal kinetic energy, which is expressed in terms of the conjugate momentum τ_α of ξ_α and the reduced masses $\mu_\alpha = m\frac{\alpha}{\alpha+1}$. As $[H_{CM}, H_{int}] = 0$, the eigenstate ψ of H can be build as a product of the form

$$\psi(\mathbf{r}_1, \ldots, \mathbf{r}_N) = \Gamma(\mathbf{R}) \; \psi_{int}(\xi_1, \ldots, \xi_{N-1}), \tag{3}$$

where Γ and ψ_{int} are defined by the equations

$$\begin{aligned} -\frac{\hbar^2}{2M}\Delta_{\mathbf{R}}\Gamma &= E_{cm}\Gamma \, , \qquad (4) \\ H_{int}\psi_{int} &= E_{int}\psi_{int} \, . \qquad (5) \end{aligned}$$

$\Gamma(\mathbf{R})$ is the c.m. wavefunction, that describes the motion of the isolated system as a whole in any inertial frame of reference. Since Γ is solution of the free Schrödinger equation, $\Gamma(\mathbf{R})$ should be an arbitrary stationary plane wave, i.e. infinitely spread and not normalizable. This leads to delocalization of $\mathbf{R}$ and arbitrary c.m. energy $E_{CM} = \hbar^2\mathbf{K}^2/(2M)$. This does not correspond to experimental situations, where the system is no longer isolated: interactions with other systems of the experimental apparatus localize the c.m.. But the formal decoupling between the c.m. motion and internal properties permits us to "let" the c.m. motion to the choice of experimental conditions, internal properties described by ψ_{int} being fully comparable to the experimental ones. Note that ψ_{int} is by definition always normalizable for the ground state of a self-bound system. The *internal* density associated to ψ_{int} is [15, 22, 23]

$$\rho_{int}(\mathbf{r}) \;=\; N\Big(\frac{N}{N-1}\Big)^3 \int d\xi_1 \cdots d\xi_{N-2} \, \big|\psi_{int}(\xi_1, \ldots, \xi_{N-2}, \tfrac{N\mathbf{r}}{N-1})\big|^2$$

$$= \; N \int d\mathbf{r}_1 \cdots d\mathbf{r}_N \; \delta(\mathbf{R}) |\psi_{int}(\mathbf{r}_1, \ldots, \mathbf{r}_N)|^2 \, \delta\big(\mathbf{r} - (\mathbf{r}_i - \mathbf{R})\big). \tag{6}$$

The density $\rho_{int}(\mathbf{r})$ is normalized to N and $\mathbf{r}$ is defined in the c.m. frame (see the delta relation in the previous equation) [1]. Note that, even if ψ_{int} can be written as a function of the $(N-1)$ Jacobi coordinates only, it can also be written as a function of the N coordinates $\mathbf{r}_i$, but one of them would be redundant [24], which is expressed by the $\delta(\mathbf{R})$ in the previous equation.

The stationary internal DFT theorem, demonstrated by various ways in Refs. [15, 25, 26], states that for a non-degenerate ground state and for given Fermion or Boson kind of particles (i.e. a given interaction u), ψ_{int} can be expressed as a unique functional of ρ_{int}, i.e. $\psi_{int}[\rho_{int}]$. As a consquence, the ground state internal energy of a self-bound system $E_{int} = (\psi_{int}[\rho_{int}]|H_{int}|\psi_{int}[\rho_{int}])$ can also be expressed as a unique functional of ρ_{int}.

A practical way to compute ρ_{int} is given by the *internal* KS scheme, developped in Ref. [15]. To set up this scheme, we assume that there exists, *in the c.m. frame*, a local single-particle potential (i.e. a N-body non-interacting system) that can reproduce exactly the density ρ_{int} of the interacting system. We develop ρ_{int} in the corresponding basis $\{\varphi^i_{int}, i = 1..N\}$ of one-body orbitals expressed in c.m. frame

$$\rho_{int}(\mathbf{r}) = \sum_{i=1}^{N} \left|\varphi^i_{int}(\mathbf{r})\right|^2 . \tag{7}$$

We refer the reader to Ref. [17], §III.C, for a justification of the introduction of N orbitals in the KS scheme, even if only $(N-1)$ coordinates are sufficient to describe internal properties. (We implicitely supposed that the particles are Fermions, but a KS scheme to describe Boson condensates can be set up similarly, equalling all the φ^i_{int}.)

The KS assumption implies $\varphi^i_{int}[\rho_{int}]$ [21]; hence, we can rewrite E_{int} as [15]

$$E_{int}[\rho_{int}] \;=\; \sum_{i=1}^{N} (\varphi^i_{int}|\frac{\mathbf{p}^2}{2m}|\varphi^i_{int}) + E_H[\rho_{int}] + E_{XC}[\rho_{int}] + \int d\mathbf{r} \; v_{int}(\mathbf{r}) \, \rho_{int}(\mathbf{r}), \tag{8}$$

where we have introduced the "non-interacting" kinetic energy $\sum_{i=1}^{N} (\varphi^i_{int}|\frac{\mathbf{p}^2}{2m}|\varphi^i_{int})$, the internal Hartree energy $E_H[\rho_{int}] = \frac{1}{2} \int d\mathbf{r} \, d\mathbf{r}' \, \rho_{int}(\mathbf{r}) \, \rho_{int}(\mathbf{r}') \, u(\mathbf{r} - \mathbf{r}')$ and the exchange-correlations energy functional

$$E_{XC}[\rho_{int}] \;=\; E_{potXC}[\rho_{int}] + E_{\Delta kin}[\rho_{int}], \tag{9}$$

composed of

$$\begin{aligned} E_{potXC}[\rho_{int}] \;&=\; \frac{1}{2} \int d\mathbf{r} \, d\mathbf{r}' \, \gamma_{int}(\mathbf{r}, \mathbf{r}') \, u(\mathbf{r} - \mathbf{r}') - E_H[\rho_{int}], \\ E_{\Delta kin}[\rho_{int}] \;&=\; (\psi_{int}| \sum_{\alpha=1}^{N-1} \frac{\tau_\alpha^2}{2\mu_\alpha} |\psi_{int}) - \sum_{i=1}^{N} (\varphi^i_{int}|\frac{\mathbf{p}^2}{2m}|\varphi^i_{int}) && (10) \\ &=\; \int d\mathbf{r}_1 \cdots d\mathbf{r}_N \delta(\mathbf{R}) \, \psi^*_{int}(\mathbf{r}_1, \ldots, \mathbf{r}_N) \sum_{i=1}^{N} \frac{\mathbf{p}_i^2}{2m} \psi_{int}(\mathbf{r}_1, \ldots, \mathbf{r}_N) - \sum_{i=1}^{N} (\varphi^i_{int}|\frac{\mathbf{p}^2}{2m}|\varphi^i_{int}). && (11) \end{aligned}$$

E_{potXC} contains the quantum exchange and correlations effets linked to the interaction u ($\gamma_{int}(\mathbf{r}, \mathbf{r}')$ is the local part of the 2-body *internal* density matrix defined in Ref. [15]; it is trivially a functional of ρ_{int}).

In the following, we call "interacting" kinetic energy the kinetic energy of the self-bound system, i.e. $\int d\mathbf{r}_1 \cdots d\mathbf{r}_N \delta(\mathbf{R}) \, \psi^*_{int}(\mathbf{r}_1, \ldots, \mathbf{r}_N) \sum_{i=1}^{N} \frac{\mathbf{p}_i^2}{2m} \psi_{int}(\mathbf{r}_1, \ldots, \mathbf{r}_N)$, and "non-interacting" kinetic energy the kinetic energy of the KS system, i.e. $\sum_{i=1}^{N} (\varphi^i_{int}|\frac{\mathbf{p}^2}{2m}|\varphi^i_{int})$. We see from Eq. (11) that $E_{\Delta kin}$ contains the exchange and "standard" correlations [2], but also the c.m. correlations (due to $\delta(\mathbf{R})$), contained in the "interacting" kinetic energy term. It is the only term of the functional that contains *explicitely* the c.m. correlations and represents the main difference with traditional DFT. Varying $E_{int}[\rho_{int}]$, Eq. (8), with respect to φ_{int}^{i*}, and imposing normality of the $\{\varphi_{int}^{i}\}$ leads to "internal" KS equations

---

[1] More generally, we can introduce a $\delta(\mathbf{R} - \mathbf{a})$ where $\mathbf{a}$ is an arbitrary translation vector, which would lead to perfectly equivalent results. We chose $\mathbf{a} = \mathbf{0}$ for simplicity, so that the formalism is formulated in the c.m. frame.

[2] "Standard" correlations means all the correlations except the c.m. correlations in the following.

$$\Big(-\frac{\hbar^2}{2m}\Delta + U_H[\rho_{int}] + U_{XC}[\rho_{int}] + v_{int}\Big)\varphi_{int}^{i} = \epsilon_i \varphi_{int}^{i}, \tag{12}$$

with $U_H[\rho_{int}](\mathbf{r}) = \delta E_H[\rho_{int}]/\delta\rho_{int}(\mathbf{r})$ and $U_{XC}[\rho_{int}](\mathbf{r}) = \delta E_{XC}[\rho_{int}]/\delta\rho_{int}(\mathbf{r})$, which are local as expected. Equations (12) have the same form as the traditional KS equations formulated for non-translationally invariant Hamiltonians [19], but we have justified their use in the c.m. frame for self-bound systems described with translational-invariant Hamiltonians, and shown that the functional form of $U_{XC}[\rho_{int}]$ differs by the inclusion of c.m. correlations [15].

We see from Eq. (11) that one has to be cautious with the meaning that is given to the non-interacting kinetic energy in mean-field like calculations. Indeed, the non-interacting kinetic energy cannot be considered as a first order approximation of the interacting kinetic energy in the general case. The difference is equal to $E_{\Delta kin}$, that can be large when c.m. correlations effects are strong, i.e. for small and intermediary size self-bound systems. For large self-bound systems $E_{\Delta kin}$ decreases (in relative value), so that the non-interacting and interacting kinetic energies become closer.

The internal DFT formalism has been generalized to time-dependent self-bound systems in Ref. [16], for instance for the description of the collision of two nuclei or laser irradiation, and to multicomponent self-bound systems in Ref. [17], for the description of self-bound systems composed by different kinds of particles (atomic nuclei, mix of ^{3}He and ^{4}He droplets, and molecular systems where the nuclei are treated explicitely). This last work permits to recover the traditional DFT formalism when one kind of particles is much heavier than the others [17], underlining why traditional DFT is essentially suited for the description of electrons (only) in molecular systems, but not for the description of self-bound systems.

B. A proposed method to obtain a c.m. correlations functional.

We first consider a system of identical Fermions (result for bosonic systems will be derived in §III F). We mentioned that the $E_{\Delta kin}[\rho_{int}]$ term defined in Eq. (10) contains the quantum exchange and "standard" correlations (remind footnote 2) contained in the interacting kinetic term, denoted $E_{\Delta kinXC}$, but also the c.m. correlations. We thus split $E_{\Delta kin}$ as

$$E_{\Delta kin} = E_{\Delta kinXC} + E_{cm}, \tag{13}$$

where the pure c.m. correlations energy E_{cm} is to be defined. Using this definition, the total exchange-"standard" correlations energy functional E_{XC}, defined by Eq. (9), can be rewritten

$$E_{XC} = E_{XC}^{stand} + E_{cm},$$

where the so-called "standard" exchange-correlations total energy is defined as $E_{XC}^{stand} = E_{potXC} + E_{\Delta kinXC}$. This term is mostly described by the parametrized functionals commonly used for mean-field like calculations of self-bound systems (see Refs. [2, 9] for a description of functionals used for nuclear systems and Ref. [27] for a description of functionals used for Helium droplets systems). But the c.m. correlations effects (described by E_{cm}) are by construction mostly *not* taken into account in those functionals (except through a renormalization of the mass in the kinetic energy term), which can affect the results, see Ref. [11]. Our goal is to set up a general well founded form for E_{cm}, which can be used for the description of all self-bound systems by simple addition to the common functionals (which, rigorously, implies a refitting of those functionals), and is numerically manageable. The idea is to start from the definition (10) of $E_{\Delta kin}$ and neglect all the exchange and "standard" correlations terms. Then, by definition (13), we will be left with E_{cm}.

In the following, we propose to search a form for E_{cm} that is functional of the KS orbitals φ_{int}^{i}. We follow this way because it allows more flexibility, while being fully coherent with DFT. Indeed, the KS orbitals are functionals of the internal density, i.e. $\varphi_{int}^{i}[\rho_{int}]$, as soon as they satisfy KS equations [21], which is the spirit of the Optimized Effective Potential (OEP) method [28–30].

C. The commonly used form for the c.m. correlations functional.

We rewrite $E_{\Delta kin}[\rho_{int}]$, Eq. (10), in the following perfectly equivalent way

$$E_{\Delta kin}[\rho_{int}] \tag{14}$$

$$
\begin{aligned}
&= \int d\mathbf{R}\delta(\mathbf{R})(\psi_{int}|\sum_{\alpha=1}^{N-1}\frac{\tau_\alpha^2}{2\mu_\alpha}|\psi_{int}) - \sum_{i=1}^{N}(\varphi^i_{int}|\frac{\mathbf{p}^2}{2m}|\varphi^i_{int}) \\
&= \int d\mathbf{R}d\xi_1\cdots d\xi_{N-1}\left(\sqrt{\delta(\mathbf{R})}\psi_{int}(\xi_1,\cdots,\xi_{N-1})\right)^*\sum_{\alpha=1}^{N-1}\frac{\tau_\alpha^2}{2\mu_\alpha}\left(\sqrt{\delta(\mathbf{R})}\psi_{int}(\xi_1,\cdots,\xi_{N-1})\right) - \sum_{i=1}^{N}(\varphi^i_{int}|\frac{\mathbf{p}^2}{2m}|\varphi^i_{int}) \\
&= \int d\mathbf{r}_1\cdots d\mathbf{r}_N\left(\sqrt{\delta(\mathbf{R})}\psi_{int}(\mathbf{r}_1,\ldots,\mathbf{r}_N)\right)^*\left(\sum_{i=1}^{N}\frac{\mathbf{p}_i^2}{2m}-\frac{\mathbf{P}^2}{2mN}\right)\left(\sqrt{\delta(\mathbf{R})}\psi_{int}(\mathbf{r}_1,\ldots,\mathbf{r}_N)\right) - \sum_{i=1}^{N}(\varphi^i_{int}|\frac{\mathbf{p}^2}{2m}|\varphi^i_{int}).
\end{aligned}
$$

$\sqrt{\delta(\mathbf{R})}\psi_{int}(\mathbf{r}_1,\ldots,\mathbf{r}_N)$ is interpreted as the c.m. frame N-body ”wavefunction” (remind that ψ_{int} has the dimension of a $(N-1)$-body wavefunction, see Eqs. (3) and (6)). This ”wavefunction” is obviously not translationally invariant (the $\delta(\mathbf{R})$ fixes the c.m. in position space and amounts to move in the c.m. frame) and is antisymmetric under the exchange of two particles (as ψ_{int} is antisymmetric). It is non null only for the $\{\mathbf{r}_i\}$ that satisfy $\mathbf{R}=\sum_{i=1}^{N}\mathbf{r}_i=0$, so that the $\{\mathbf{r}_i\}$ become the c.m. frame coordinates.

Within the internal DFT formalism, the commonly used approximation to treat the c.m. correlations can be recovered by supposing that the KS Slater Determinant, denoted ψ^{aux}, is a good first order approximation of the c.m. frame N-body ”wavefunction”:

$$
\sqrt{\delta(\mathbf{R})}\psi_{int}(\mathbf{r}_1,\ldots,\mathbf{r}_N) \quad\approx\quad \psi^{aux}(\mathbf{r}_1,\ldots,\mathbf{r}_N), \tag{15}
$$

where (P are the possible permutations of the coordinates and p the number of transpositions of P)

$$
\psi^{aux}(\mathbf{r}_1,\ldots,\mathbf{r}_N)=\frac{1}{\sqrt{N!}}\sum_{P}(-1)^p\Pi_{i=1}^{N}\varphi_{int}^{P(i)}(\mathbf{r}_i).
$$

Inserting this approximation in (14) and following the method described in §II B (the ”standard” correlations are by construction neglected and the exchange terms cancel naturally), we obtain

$$
\begin{aligned}
&E_{\Delta kin}\rightarrow E_{cm}[\{\varphi^k_{int}\}] \\
&\quad= -\left(\psi^{aux}\middle|\frac{\mathbf{P}^2}{2mN}\middle|\psi^{aux}\right) \\
&\quad= -\frac{\hbar^2}{2mN}\sum_{i=1}^{N}\int d\mathbf{r}\;\varphi^{i*}_{int}(\mathbf{r})\Delta_{\mathbf{r}}\varphi^i_{int}(\mathbf{r}) - \frac{\hbar^2}{2mN}\sum_{i,j=1}^{N}\int d\mathbf{r}\;\varphi^{i*}_{int}(\mathbf{r})\nabla_{\mathbf{r}}\varphi^i_{int}(\mathbf{r})\int d\mathbf{r}'\;\varphi^{j*}_{int}(\mathbf{r}')\nabla_{\mathbf{r}'}\varphi^j_{int}(\mathbf{r}').
\end{aligned}
\tag{16}
$$

We recover the commonly used form for the c.m. correlations functional (note that, in practise, the second term of the last line of Eq. (16) is often neglected to reduce the numerical cost) [11].

The internal DFT formalism permits us to shed a new light on the validity of the approximation (16). It holds if and only if the approximation (15) holds at least to first order. But this cannot be the case in general because ψ^{aux} is far form being null when $\sum_{i=1}^{N}\mathbf{r}_i\neq 0$. Moreover, ψ^{aux} contains a c.m. vibration, typical of Slater determinants (i.e. $(\psi^{aux}|\mathbf{P}^n|\psi^{aux})\neq 0$ for $n\geq 2$) [1, 2], whereas $\sqrt{\delta(\mathbf{R})}\psi_{int}$ does not contain such a vibration (i.e. $(\psi^{int}|\mathbf{P}^n|\psi^{int})=0$, $\forall n$). Thus, we cannot expect for a systematic satisfying improvement with this form [11].

In the next section, we propose a better founded form for the c.m. correlations energy functional, where the c.m. correlations (the $\delta(\mathbf{R})$ term) appear explicitely.

III. A GENERAL NEW FORM FOR A LOCAL C.M. CORRELATIONS POTENTIAL.

A. The idea.

We adopt a different point of view than the one of §II C. We start from $E_{\Delta kin}[\rho_{int}]$ written as in Eq. (11) (instead of (14)) and pose (instead of (15))

$$
\psi_{int}(\mathbf{r}_1,\ldots,\mathbf{r}_N) \quad\rightarrow\quad \frac{1}{\Gamma^{aux}(\mathbf{R})}\psi^{aux}(\mathbf{r}_1,\ldots,\mathbf{r}_N). \tag{17}
$$

The reasons of the introduction of the 1-body ”wave-function” $\Gamma^{aux}(\mathbf{R})$ (that implicitely depends on the number of particles N) are the followings:

- ψ_{int} has the dimension of a $(N-1)$-body wavefunction, whereas the KS Slater Determinant $\psi^{aux}(\mathbf{r}_1,\ldots,\mathbf{r}_N)$ has the dimension of a N-body wavefunction. Dividing ψ^{aux} by Γ^{aux} permits to recover the dimension of a $(N-1)$-body wavefunction while preserving antisymmetry.
- The KS Slater determinant ψ^{aux} contains a c.m. vibration, as already mentioned. This is not a problem from the KS point of view, because all is wanted is that ψ^{aux} reproduces the correct ρ_{int}, the KS Slater determinant representing nothing more than an auxiliary. Hovever, if we wan't to use ψ^{aux} to approximate ψ_{int}, the c.m. vibration of ψ^{aux} has to be "substracted", because ψ_{int} should not contain such a vibration. Γ^{aux} represents the proposed way to "substract" the c.m. vibration contained in ψ^{aux}.

  In the particular case of the harmonic oscillator (i.e. when the interaction u is parabolic), we can always achieve the separation $\psi^{aux}(\mathbf{r}_1,\ldots,\mathbf{r}_N)=\Gamma^{aux}(\mathbf{R})\times F(\xi_1,\ldots,\xi_{N-1})$ [13]. Thus, the use of the form (17) permits to "substract" exactly and directly all the c.m. vibration contained in ψ^{aux}.

  In the general case, $\psi^{aux}(\mathbf{r}_1,\ldots,\mathbf{r}_N)$ cannot be separated into $\Gamma^{aux}(\mathbf{R})\times F(\xi_1,\ldots,\xi_{N-1})$. However, all we wan't is that the form of Γ^{aux} would allow sufficient flexibility to lead to a satisfying result for both E_{cm} and ρ_{int} (then, even if the "substraction" is not "direct" because there is no separation of the c.m. motion in ψ^{aux}, it is "indirect" because it leads to the correct final result).
- As we will see, the final result has a clear Physical meaning, which shows the pertinence of our proposition.
- The $\delta(\mathbf{R})$ contained in (11) breaks translational invariance, so that even if $\frac{1}{\Gamma^{aux}(\mathbf{R})}\psi^{aux}(\mathbf{r}_1,\ldots,\mathbf{r}_N)$ is not translationally invariant (contrary to ψ_{int}), the final result won't be affected if a good form for Γ^{aux} is found.

Necessary properties that Γ^{aux} should satisfy will be discussed in §III D.

B. The result.

We first note that, when inserted in (11), the approximation (17) does not necessarilly lead to a real $E_{\Delta kin}$ [3]. To overcome this problem (which would affect time-dependent results), we insert (17) in $E_{\Delta kin}$ and keep only the real part. We then obtain an approximation of the exact $E_{\Delta kin}$ where the "standard" correlations have been neglected by construction. As discussed in §II B, it remains to neglect the exchange terms in the result to obtain E_{cm}. The calculation is detailled in Appendix A. The final result is

$$\begin{aligned}
&E_{\Delta kin}\rightarrow E_{cm}[\{\varphi^k_{int}\}]=\\
&\quad-\frac{\hbar^2}{2m}\sum_{i=1}^{N}\int d\mathbf{r}\ \varphi^{i*}_{int}(\mathbf{r})\Delta_{\mathbf{r}}\varphi^{i}_{int}(\mathbf{r})\times\Big(\frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}\int d\mathbf{r}'\ |\varphi^{l\neq i}_{int}(\mathbf{r}')|^2\times f_{i,l\neq i}[\{\varphi^{k\neq i,l}_{int}\}](\mathbf{r}+\mathbf{r}')-1\Big)\\
&\quad-\frac{\hbar^2}{2mN}\ \Re e\Big(\frac{1}{\Gamma^{aux*}(\mathbf{0})}\Delta_{\mathbf{R}}\frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}}\Big)\times\int d\mathbf{r}\ |\varphi^{i}_{int}(\mathbf{r})|^2\int d\mathbf{r}'\ |\varphi^{l\neq i}_{int}(\mathbf{r}')|^2\times f_{i,l\neq i}[\{\varphi^{k\neq i,l}_{int}\}](\mathbf{r}+\mathbf{r}')\\
&\quad+PureImaginary[\{\varphi^k_{int}\}],
\end{aligned}\tag{18}$$

where the functional $PureImaginary$ is defined as

$$PureImaginary[\{\varphi^k_{int}\}]=i\frac{\hbar^2}{2m}\Im m\sum_{i=1}^{N}\int d\mathbf{r}\ \varphi^{i*}_{int}(\mathbf{r})\Delta_{\mathbf{r}}\varphi^{i}_{int}(\mathbf{r})\times\frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}\int d\mathbf{r}'\ |\varphi^{l\neq i}_{int}(\mathbf{r}')|^2\times f_{i,l\neq i}[\{\varphi^{k\neq i,l}_{int}\}](\mathbf{r}+\mathbf{r}').\tag{19}$$

It counter acts the pure imaginary part of the first line of (18) and becomes null in the real (stationary) case. We keep it for generality (time-dependent case).

The "two particles c.m. correlations functional" is defined as (D is the dimension in which the calculations are done: D=1,2 or 3)

$$f_{i,l\neq i}[\{\varphi^{k\neq i,l}_{int}\}](\tilde{\mathbf{r}})=N^D\int\Pi^N_{\substack{j=1\\ j\neq i,l}}d\mathbf{r}_j\delta\Big(\sum_{\substack{k=1\\ k\neq i,l}}^{N}\mathbf{r}_k+\tilde{\mathbf{r}}\Big)\Pi^N_{\substack{j=1\\ j\neq i,l}}|\varphi^{j}_{int}(\mathbf{r}_j)|^2.\tag{20}$$

---

[3] This is fundamentally due to the fact that, in the general case, the form (17) cannot be rewritten as a function of the $\{\xi_\alpha\}$ only.

$f_{i,l\neq i}$ couples the particles i and l together in (18). The meaning and properties of this functional will be detailed in §III C.

The potentials U^l_{cm} corresponding to E_{cm} are defined by [4]

$$\begin{aligned}
U^l_{cm}(\mathbf{r})\varphi^l_{int}(\mathbf{r}) &= \frac{\delta E_{cm}[\{\varphi^k_{int}\}]}{\delta\varphi^{l*}_{int}(\mathbf{r})} \\
&= -\frac{\hbar^2}{2m}\Big\{\Delta_{\mathbf{r}}\varphi^l_{int}(\mathbf{r}) \times \Big(\frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}\int d\mathbf{r}'|\varphi^{m\neq l}_{int}(\mathbf{r}')|^2 f_{l,m\neq l}[\{\varphi^{k\neq l,m}_{int}\}](\mathbf{r}+\mathbf{r}') - 1\Big) \\
&\qquad + \frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}\varphi^l_{int}(\mathbf{r})\sum_{\substack{i=1\\ i\neq l}}^{N}\int d\mathbf{r}'\ \varphi^{i*}_{int}(\mathbf{r}')\Delta_{\mathbf{r}'}\varphi^i_{int}(\mathbf{r}') \times f_{i,l\neq i}[\{\varphi^{k\neq i,l}_{int}\}](\mathbf{r}+\mathbf{r}')\Big\} \\
&\quad -\frac{\hbar^2}{2mN}\ \Re e\Big(\frac{1}{\Gamma^{aux*}(\mathbf{0})}\Delta_{\mathbf{R}}\frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}}\Big)\times\varphi^l_{int}(\mathbf{r})\int d\mathbf{r}'|\varphi^{m\neq l}_{int}(\mathbf{r}')|^2 \times f_{l,m\neq l}[\{\varphi^{k\neq l,m}_{int}\}](\mathbf{r}+\mathbf{r}') \\
&\quad + \frac{\delta}{\delta\varphi^{l*}_{int}(\mathbf{r})}PureImaginary[\{\varphi^k_{int}\}], \qquad (21)
\end{aligned}$$

where the last line is obviously null in the real (stationary) case. Note that the potentials U^l_{cm} are not common for all states. This is due to the fact that $E_{cm}[\{\varphi^k_{int}\}]$ is orbital dependent, which necessits extra measures to recover a common potential or, equivalently, to preserve orthonormalization. A way to overcome this problem is to use the OEP method, which permits us to find the potential common to all states that reproduces the most accurately the effect of the U^l_{cm} potentials. We refer the reader to Refs. [28–30] for the exhaustive equations. As the full OEP result is very costly numerically, it is often simplified. Popular is, e.g., the Krieger-Li-Iafrate (KLI) approach and, in a further step of simplification, the Slater approximation [31, 32]. As our goal is to find a numerically cheap form for the local c.m. potential, we detail thereafter only the Slater approximation. It gives:

$$U^{Slat}_{cm}(\mathbf{r}) = \sum_{i=l}^{N}\frac{|\varphi^l_{int}(\mathbf{r})|^2}{\rho_{int}(\mathbf{r})}U^l_{cm}(\mathbf{r}). \qquad (22)$$

C. Properties of $f_{i,l\neq i}$.

The definition (20) of the two particles c.m. correlations functional $f_{i,l\neq i}$ shows that:

1. it is real and has the dimension of a density,
2. it is normalized to N^D, i.e. $\int d\tilde{\mathbf{r}} f_{i,l\neq i}(\tilde{\mathbf{r}}) = N^D$,
3. $\lim_{\tilde{\mathbf{r}}\to\pm\infty} f_{i,l\neq i}(\tilde{\mathbf{r}}) = 0$,
4. it is a "multiconvolution" of all single densities, unless these associated to orbitals i and l.

The first three points permit to explicit the physical meaning of $\frac{1}{N^D}f_{i,l\neq i}(\mathbf{r}+\mathbf{r}')$: it is the probability that particle $l\neq i$ has position $\mathbf{r}'$, given particle i has position $\mathbf{r}$. Indeed, because of the c.m. correlations, the positions of those particles are not independent and are coupled by $f_{i,l\neq i}$. As a consequence, this coupling appears in the c.m. correlations energy (18) and potentials (21): because of the c.m. correlations, every single orbital φ^i_{int} is coupled to every single orbitals $\varphi^{l\neq i}_{int}$ through $f_{i,l\neq i}$.

To understand better this coupling, note that $f_{i,l\neq i}$ can be rewritten

$$\begin{aligned}
f_{i,l\neq i}[\{\varphi^{k\neq i,l}_{int}\}](\tilde{\mathbf{r}}) &= 2^D\delta(\tilde{\mathbf{r}}) \quad if \quad N=2, \\
& 3^D|\varphi^{k\neq i,l}_{int}(-\tilde{\mathbf{r}})|^2 \quad if \quad N=3,
\end{aligned}$$

---

[4] A usefull and straightforward relation to achieve the variation is ($\delta_{l,i}$ is the Kronecker symbol): $\frac{\delta}{\delta\varphi^{l*}_{int}(\mathbf{r})}\int d\mathbf{r}'\ |\varphi^{m\neq i}_{int}(\mathbf{r}')|^2 \times f_{i,m\neq i}[\{\varphi^{k\neq i,m}_{int}\}](\mathbf{r}'+\mathbf{r}'') = (1-\delta_{l,i})\ \varphi^l_{int}(\mathbf{r})\ f_{i,l\neq i}[\{\varphi^{k\neq i,l}_{int}\}](\mathbf{r}+\mathbf{r}'')$.

$$N^D \int \Pi^N_{\substack{j=1\\ j\neq i,l,m}} d\mathbf{r}_j \Pi^N_{\substack{j=1\\ j\neq i,l,m}} |\varphi^j_{int}(\mathbf{r}_j)|^2 |\varphi^m_{int}\Big(-\sum_{\substack{k=1\\ k\neq i,l,m}}^{N} \mathbf{r}_k - \tilde{\mathbf{r}}\Big)|^2 \quad if \quad N \geq 4, \tag{23}$$

$$\ldots$$

$$Constant, \quad for\ very\ large\ N\ (limit\ of\ a\ Fermi\ Gaz).$$

We see that, in the 2 particles case, $f_{1,2}$ is proportional to a the steep delta function. Thus, if particle 1 has position $\mathbf{r}$, particle 2 will have position **-r**, so that the c.m. remains stuck to $\mathbf{R} = \mathbf{0}$. In the 3 particles case, $f_{i,l\neq i}$ has a larger width, because the introduction of a third particle allows more freedom to the motion of the 2 other particles, while preserving $\mathbf{R} = \mathbf{0}$. As $f_{i,l\neq i}$ is a multiconvolution of all the single densities unless the ones associated to orbitals i and l, it should fastly become very little dependent of i and l as N grows. For $N \geq 4$, the width of $f_{i,l\neq i}$ will increase as N grows, because of the multiconvolution form of $f_{i,l\neq i}$. Indeed, a larger number of particles allows more liberty to the motion of 2 of these particles, while preserving $\mathbf{R} = \mathbf{0}$. For very large N, the system tends to a Fermi gaz, so that $f_{i,l\neq i}$ tends to become constant and delocalized in the whole space, i.e. the motions the particles tend to become independent. The c.m. correlations can then be neglected, as expected.

Practically speaking, we see that the numerical cost of the whole scheme lies in the calculation of $f_{i,l\neq i}$ for $N \geq 4$, i.e. the calculation of the multiconvolution of Eq. (23). At first sight, it seems to be discrimatory for large N. But a mathematical property of the convolutions under Fourier transforms makes it manageable. In Appendix B is demonstrated the so called "multiconvolution theorem". Its direct application to $f_{i,l\neq i}$ for $N \geq 4$ gives ($\mathcal{T}$ denotes the Fourier transform as defined in Appendix B, Eq. (B1))

$$f_{i,l\neq i}[\{\varphi^{k\neq i,l}_{int}\}](\tilde{\mathbf{r}}) = N^D \times \mathcal{T}^{-1}\Big[\Pi^N_{\substack{k=1\\ k\neq i,l}} \mathcal{T}[|\varphi^k_{int}|^2]\Big](-\tilde{\mathbf{r}}).$$

This permits to fasten drastically the numerical calculation of $f_{i,l\neq i}$, which becomes manageable even for large systems. Indeed, once every $\mathcal{T}[|\varphi^j_{int}|^2]$ are calculated, $f_{i,l\neq i}$ is given by the inverse Fourier transform of their direct product, so that the numerical cost of $f_{i,l\neq i}$ equals the numerical cost of $(N+1)$ Fast Fourier Transforms when $N \geq 4$.

D. Properties of Γ^{aux}.

Remind that Γ^{aux} implicitely depends on N. We see from (18) that Γ^{aux} should be two times derivable and that only the values of Γ^{aux} and of its second derivative in $\mathbf{R} = \mathbf{0}$ are of interest to reproduce E_{cm}. We now characterize those values.

First, we note that $|\Gamma^{aux}(\mathbf{0})|^2$ is imposed by the normalization condition on the approximation (17) we used for ψ_{int} [5]:

$$\begin{aligned} 1 = (\psi_{int}|\psi_{int}) &= \int d\mathbf{r}_1 \cdots d\mathbf{r}_N\ \delta(\mathbf{R}) |\psi_{int}(\mathbf{r}_1, \ldots, \mathbf{r}_N)|^2 \\ \Rightarrow \quad |\Gamma^{aux}(\mathbf{0})|^2 &= \int d\mathbf{r}\ d\mathbf{r}'\ |\varphi^i_{int}(\mathbf{r})|^2 |\varphi^{l\neq i}_{int}(\mathbf{r}')|^2 f_{i,l\neq i}[\{\varphi^{k\neq i,l}_{int}\}](\mathbf{r}+\mathbf{r}'). \end{aligned} \tag{24}$$

Numerically speaking, this condition will be satisfied self-consistently, starting from a reasonnable initial value for $|\Gamma^{aux}(\mathbf{0})|^2$ and rescaling it at every loop so that it satisfies (24).

It remains to characterize the value of $\frac{1}{\Gamma^{aux*}(\mathbf{0})} \Delta_{\mathbf{R}} \frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}}$. To that aim, we use the following results demonstrated in Appendix C:

$$\lim_{N\to+\infty} |\Gamma^{aux}(\mathbf{0})|^2 \to +\infty, \tag{25}$$

$$\lim_{N\to+\infty} \frac{1}{N} \times \frac{1}{\Gamma^{aux*}(\mathbf{0})} \Delta_{\mathbf{R}} \frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}} \times |\Gamma^{aux}(\mathbf{0})|^2 \to 0. \tag{26}$$

The most simple and natural [4] form for Γ^{aux} that meets those criteria (and is "exact" in the case where the interaction u is parabolic) is a Gaussian

$$\Gamma^{aux}(\mathbf{R}) = \Big(\frac{K(N)}{\pi}\Big)^{D/4} \exp\Big\{-\frac{K(N)}{2}\sum_{i=1}^{D} R_i^2\Big\}, \tag{27}$$

---

[5] The second line of Eq. (24) is obtained introducing the form (17) for ψ_{int} in the first line of Eq. (24) and neglecting the exchange terms.

where R_i are the coordinates of $\mathbf{R}$ in D dimensions and $K(N)$ is a two times derivable function of N. With this form

$$|\Gamma^{aux}(\mathbf{0})|^2 = \Big(\frac{K(N)}{\pi}\Big)^{D/2}, \tag{28}$$

$$\frac{1}{\Gamma^{aux*}(\mathbf{0})}\Delta_{\mathbf{R}}\frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}} = \pi^{D/2}\times D\times\Big(K(N)\Big)^{1-D/2}. \tag{29}$$

This permits us to define completely the c.m. correlation energy (18) and potential (21), without introduction of any free parameter. Indeed, as previously mentionned, $|\Gamma^{aux}(\mathbf{0})|^2$ is defined by Eq. (24) and should be rescaled at every numerical loop so that it satisfies this relation. This defines $K(N)$ at each loop by (28), which defines $\frac{1}{\Gamma^{aux*}(\mathbf{0})}\Delta_{\mathbf{R}}\frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}}$ by (29), and we are done.

With this method, there is no need to define analytically K as a function of N; $K(N)$ is obtained numerically for every given N as indicated. It would nevertheless be interesting to obtain an approximate analytical form for $K(N)$, at least to start the numerical iterations with a pertinent initial condition. To that aim, we notice that the conditions (25) and (26), together with the equalities (28) and (29), imply the following conditions on $K(N)$

$$\lim_{N\to+\infty}\Big(K(N)\Big)^{D/2} \to +\infty,$$

$$\lim_{N\to+\infty}\frac{1}{N}K(N) \to 0. \tag{30}$$

A straightforward form for $K(N)$ that satisfies those two constraints is

$$K(N) = A\times N^a \quad , \quad \textit{where } 0 < a < 1. \tag{31}$$

The corresponding energy associated to Γ^{aux} is then

$$E_{\Gamma^{aux}} = -\frac{\hbar^2}{2Nm}(\Gamma^{aux}|\Delta_{\mathbf{R}}|\Gamma^{aux}) = \frac{\hbar^2}{2Nm}\frac{D}{4}K(N) = \frac{\hbar^2}{2m}\frac{D}{4}A\times N^{a-1}.$$

$E_{\Gamma^{aux}}$ is then proportional to N^{a-1}. Even if $E_{\Gamma^{aux}}$ has, strictly speaking, no Physical meaning, is is reasonnable to assume that its variation according to N should approximately be proportional to the variation of the energy associated to the c.m. vibration obtained in mean-field like calculation, see Ref. [11]. For the nuclear case, the c.m. vibration energy evaluated for 3D harmonic oscillator states is proportional to $N^{-1/3}$; this variation can be reproduced with $a = 2/3$. The c.m. correlation energy evaluated with a *a posteriori* fit with 3D mean-field like calculations is proportional to $N^{-0.2}$; this variation can be reproduced with $a \approx 0.8$ [11]. This is fully coherent with the fact that $0 < a < 1$, as deduced by our previous considerations, which reinforces the pertinence of the Gaussian form, that we will use in the following numerical calculations. However, the search for other forms, i.e. with other variations around $\mathbf{R} = \mathbf{0}$, should be continued to obtain the most precise description of self-bound systems in fully realistic calculations.

E. An explicit density functional for the c.m correlations energy.

The functional proposed in §III B is well suited for stationary calculations, but not for time-dependent ones, because the Slater approximation, as well as the KLI approximation, do not permit to preserve energy conservation (as those approximations are not perfectly variational [33]). Only the full time-dependent OEP result [34] will achieve energy conservation, but at the price of a much larger numerical cost. It thus would be interesting to find a functional that overcomes those pathologies in the time-dependent case.

In this section, we propose a further step of approximation that will allow to obtain an explicit functional of ρ_{int} for the c.m. correlations energy. We will do the Local Density Approximation (LDA) on the result of §III B, which consists in doing the approximation that the system is *locally* homogeneous [21, 35]. Despite of its simplicity, this approximation has proven to be very satisfying for the description of a wide range of systems and not only large ones [2, 21]. To do the LDA, we first do the replacement

$$|\varphi^l_{int}(\mathbf{r})|^2 \ \to\ \frac{1}{N}\rho_{int}(\mathbf{r}) \tag{32}$$

everywhere the single density terms $|\varphi^l_{int}|^2$ appear into E_{cm}, Eq. (18). In the obtained functional, the only remaining term that is not an explicit functional of ρ_{int} is $\sum_{i=1}^N \varphi^{i*}_{int}(\mathbf{r})\Delta_{\mathbf{r}}\varphi^i_{int}(\mathbf{r})$. As we consider a system composed of Fermions, we do the Thomas-Fermi approximation [21], i.e. the replacement

$$\sum_{i=1}^{N} \varphi^{i*}_{int}(\mathbf{r})\Delta_{\mathbf{r}}\varphi^{i}_{int}(\mathbf{r}) \;\rightarrow\; -\frac{3}{5}C\rho^{5/3}_{int}(\mathbf{r}), \tag{33}$$

where $C = (\frac{3\pi^2}{\gamma})^{2/3}$ and γ is the degeneracy [6]. We obtain as a final result the c.m. correlations energy written as an explicit functional of ρ_{int}:

$$\begin{aligned} E^{LDA}_{cm}[\rho_{int}] \;=\; & \frac{\hbar^2}{2m}\int d\mathbf{r}\;\frac{3}{5}C\rho^{5/3}_{int}(\mathbf{r})\;\Big(\frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}\int d\mathbf{r}'\;\frac{1}{N}\rho_{int}(\mathbf{r}')\times f_2[\rho_{int}](\mathbf{r}+\mathbf{r}')-1\Big) \\ & -\frac{\hbar^2}{2mN}\;\Re e\Big(\frac{1}{\Gamma^{aux*}(\mathbf{0})}\Delta_{\mathbf{R}}\frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}}\Big)\times\int d\mathbf{r}\;\frac{1}{N}\rho_{int}(\mathbf{r})\int d\mathbf{r}'\;\frac{1}{N}\rho_{int}(\mathbf{r}')\times f_2[\rho_{int}](\mathbf{r}+\mathbf{r}'), \end{aligned} \tag{34}$$

where the "two particles *average* c.m. correlations functional" is defined as

$$f_2[\rho_{int}](\tilde{\mathbf{r}}) = N^D\frac{1}{N^{N-2}}\int d\mathbf{r}_1\ldots d\mathbf{r}_{N-2}\;\rho_{int}(\mathbf{r}_1)\times\cdots\times\rho_{int}(\mathbf{r}_{N-2})\;\delta\big(\sum_{k=1}^{N-2}\mathbf{r}_k+\tilde{\mathbf{r}}\big). \tag{35}$$

The corresponding unique c.m. correlations potential is given by [7]

$$\begin{aligned} \frac{\delta E^{LDA}_{cm}[\rho_{int}]}{\delta\rho_{int}(\mathbf{r})} = U^{LDA}_{cm}[\rho_{int}](\mathbf{r}) \;=\; & \frac{\hbar^2}{2m}\Big\{C\rho^{2/3}_{int}(\mathbf{r})\times\Big(\frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}\int d\mathbf{r}'\frac{1}{N}\rho_{int}(\mathbf{r}')f_2[\rho_{int}](\mathbf{r}+\mathbf{r}')-1\Big) \\ & +\frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}\int d\mathbf{r}'\;\frac{3}{5}C\rho^{5/3}_{int}(\mathbf{r}')\times\frac{N-1}{N}f_2[\rho_{int}](\mathbf{r}+\mathbf{r}')\Big\} \\ & -\frac{\hbar^2}{2mN}\;\Re e\Big(\frac{1}{\Gamma^{aux*}(\mathbf{0})}\Delta_{\mathbf{R}}\frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}}\Big)\times\int d\mathbf{r}'\frac{1}{N}\rho_{int}(\mathbf{r}')\times f_2[\rho_{int}](\mathbf{r}+\mathbf{r}'). \end{aligned} \tag{36}$$

f_2 has the following properties:

1. it is real and has the dimension of a density,
2. it is normalized to N^D, i.e. $\int d\tilde{\mathbf{r}} f_2(\tilde{\mathbf{r}}) = N^D$,
3. $\lim_{\tilde{\mathbf{r}}\to\pm\infty} f_2(\tilde{\mathbf{r}}) = 0$,
4. it is a "multiconvolution" of $(N-2)$ ρ_{int}.

The first three properties and the considerations of §III C permit to interpret $\frac{1}{N^D}f_2(\mathbf{r}+\mathbf{r}')$ as the average probability that one particle has position $\mathbf{r}'$, given another particle has position $\mathbf{r}$. f_2 can be rewritten more explicitely in the following way:

$$\begin{aligned} f_2[\rho_{int}](\tilde{\mathbf{r}}) \;=\; & 2^D\delta(\tilde{\mathbf{r}}) \quad if \quad N=2, \\ & 3^D\frac{1}{N}\rho_{int}\big(-\tilde{\mathbf{r}}\big) \quad if \quad N=3, \\ & N^D\frac{1}{N^{N-2}}\int d\mathbf{r}_1\ldots d\mathbf{r}_{N-3}\;\rho_{int}(\mathbf{r}_1)\times\cdots\times\rho_{int}(\mathbf{r}_{N-3})\;\rho_{int}\big(-\sum_{k=1}^{N-3}\mathbf{r}_k-\tilde{\mathbf{r}}\big) \quad if \quad N\geq 4, \\ & \ldots \\ & Constant, \quad for\ very\ large\ N\ (limit\ of\ a\ Fermi\ Gaz). \end{aligned} \tag{37}$$

---

[6] More precisely, if we take into account the dimension D in which the calculations are done, we have $\sum_{i=1}^{N}\varphi^{i*}_{int}(\mathbf{r})\Delta_{\mathbf{r}}\varphi^{i}_{int}(\mathbf{r}) \rightarrow -\frac{D}{2+D}C_D\rho^{\frac{2+D}{D}}_{int}(\mathbf{r})$, where $C_3 = (\frac{3\pi^2}{\gamma})^{2/3}$, $C_2 = \frac{4\pi}{\gamma}$ and $C_1 = \frac{2\pi^2}{\gamma}$.

[7] A usefull and easy to demonstrate relation to achieve the variation is the following: $\frac{\delta}{\delta\varphi^{l*}_{int}(\mathbf{r})}\int d\mathbf{r}'\;\frac{1}{N}\rho_{int}(\mathbf{r}')\times f_2[\rho_{int}](\mathbf{r}'+\mathbf{r}'') = \varphi^{l}_{int}(\mathbf{r})\;\frac{N-1}{N}\;f_2[\rho_{int}](\mathbf{r}+\mathbf{r}'')$.

Still, we see that the numerical cost lies in the calculation of $f_2[\rho_{int}]$ for $N \geq 4$. To reduce this cost, we use the "multiconvolution theorem" demonstrated in Appendix B. Its direct application to $f_2[\rho_{int}]$ for $N \geq 4$ gives (using the definition Eq. (B1) for the Fourier transform $\mathcal{T}$)

$$f_2[\rho_{int}](\tilde{\mathbf{r}}) = N^D \frac{1}{N^{N-2}} \times \mathcal{T}^{-1}\Big[(\mathcal{T}[\rho_{int}])^{N-2}\Big](-\tilde{\mathbf{r}}).$$

This permits to fasten drastically the numerical calculation of f_2, which becomes manageable even for very large systems. Indeed, one simply calculates $\mathcal{T}[\rho_{int}]$, puts it to power $(N-2)$ and calculates its inverse Fourier transform. Thus, the numerical cost of the calculation of f_2 is equal to two Fast Fourier Transforms for all $N \geq 4$.

Moreover, this scheme is perfectly variational, contrary to the one of §III B, as already mentioned, and thus will be suitable for stationary calculations as well as for time-dependent ones (i.e. it will achieve energy conservation if time-independent v_{int} is used).

F. And for Bosons?

The c.m. correlations energy functional for Bosons condensates is obtained simply by doing $\varphi^i_{int} \to \varphi_{int}$, thus $\rho_{int} = N|\varphi_{int}|^2$, in (18). Posing $\varphi_{int}(\mathbf{r}) = \sqrt{\rho_{int}(\mathbf{r})/N}$, we obtain

$$\begin{aligned} E_{cm}[\rho_{int}] \;=\; & -\frac{\hbar^2}{2m}\int d\mathbf{r}\,\sqrt{\rho_{int}(\mathbf{r})}\Delta_{\mathbf{r}}\sqrt{\rho_{int}(\mathbf{r})}\times\Big(\frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}\int d\mathbf{r}'\,\frac{1}{N}\rho_{int}(\mathbf{r}')\times f_2[\rho_{int}](\mathbf{r}+\mathbf{r}')-1\Big) \\ & -\frac{\hbar^2}{2mN}\,\Re e\Big(\frac{1}{\Gamma^{aux*}(\mathbf{0})}\Delta_{\mathbf{R}}\frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}}\Big)\times\int d\mathbf{r}\,\frac{1}{N}\rho_{int}(\mathbf{r})\int d\mathbf{r}'\,\frac{1}{N}\rho_{int}(\mathbf{r}')\times f_2[\rho_{int}](\mathbf{r}+\mathbf{r}'), \end{aligned} \tag{38}$$

where the two particles *average* c.m. correlations functional is defined as in Eq. (35) and (37), and the corresponding c.m. correlations potential is given by

$$\begin{aligned} \frac{\delta E_{cm}[\rho_{int}]}{\delta\rho_{int}(\mathbf{r})} = U_{cm}[\rho_{int}](\mathbf{r}) \;=\; & -\frac{\hbar^2}{2m}\Big\{\frac{1}{\sqrt{\rho_{int}(\mathbf{r})}}\Delta_{\mathbf{r}}\sqrt{\rho_{int}(\mathbf{r})}\times\Big(\frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}\int d\mathbf{r}'\frac{1}{N}\rho_{int}(\mathbf{r}')f_2[\rho_{int}](\mathbf{r}+\mathbf{r}')-1\Big) \\ & +\frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}\int d\mathbf{r}'\,\sqrt{\rho_{int}(\mathbf{r})}\Delta_{\mathbf{r}}\sqrt{\rho_{int}(\mathbf{r})}\times\frac{N-1}{N}f_2[\rho_{int}](\mathbf{r}+\mathbf{r}')\Big\} \\ & -\frac{\hbar^2}{2mN}\,\Re e\Big(\frac{1}{\Gamma^{aux*}(\mathbf{0})}\Delta_{\mathbf{R}}\frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}}\Big)\times\int d\mathbf{r}'\frac{1}{N}\rho_{int}(\mathbf{r}')\times f_2[\rho_{int}](\mathbf{r}+\mathbf{r}'). \end{aligned} \tag{39}$$

This potential is common to all states, so that no extra measures to preserve orthonormalization are to be taken, and it is strictly variational, so that it is usable in the time-dependent case.

IV. FIRST NUMERICAL RESULTS.

In the following, we will consider 1D calculations, which permit to understand better some features of the internal DFT formalism and to include more easilly various particle-particle interactions.

A. Model system composed of two different particles with a strong interaction.

1. *The model and the benchmark.*

We consider a stationary multicomponent self-bound system composed of two different particles, to maximize the c.m. correlations effects. We suppose that the two particles have the same mass m and are coupled by a strong interaction, which models for instance a proton and a neutron. The first particle has laboratory coordinate $r^{(1)}$ and the second has laboratory coordinate $r^{(2)}$. The reduced mass is $\mu = m/2$ and the Jacobi coordinate $\xi = r^{(1)} - r^{(2)}$. We suppose that the interaction between the two particles is parabolic (harmonic oscillator), so that the Hamiltonian is

$$H \;=\; \frac{{p^{(1)}}^2}{2m} + \frac{{p^{(1)}}^2}{2m} + \frac{1}{4}m\omega^2(r^{(1)} - r^{(2)})^2$$

$$= \frac{\tau^2}{2\mu} + \frac{1}{2}\mu\omega^2\xi^2. \tag{40}$$

The exact solution can be written in an analytical form (ψ_{int} should not be anti-symmetrized because we deal with two different particles)

$$\psi_{int}(\xi) = \Big(\frac{\mu\omega}{\pi\hbar}\Big)^{\frac{1}{4}} \exp\Big\{-\frac{1}{2}\frac{\mu\omega}{\hbar}\xi^2\Big\} \quad \Leftrightarrow \quad \psi_{int}(r^{(1)}-r^{(2)}) = \Big(\frac{m\omega}{2\pi\hbar}\Big)^{\frac{1}{4}} \exp\Big\{-\frac{1}{2}\frac{m\omega}{2\hbar}(r^{(1)}-r^{(2)})^2\Big\}. \tag{41}$$

The corresponding energy is $E_{int} = \frac{1}{2}\hbar\omega$ and the c.m. frame one-body densities for each kind of particle are [17] ($R = (r^{(1)} + r^{(2)})/2$):

$$\begin{aligned}
\rho^{(l)}_{int}(r) &= \int dr^{(1)}dr^{(2)}\delta(R)|\psi_{int}(r^{(1)}-r^{(2)})|^2\delta\big(r-(r^{(l)}-R)\big) \\
&= 2|\psi_{int}(2r)|^2 = \sqrt{\frac{2m\omega}{\pi\hbar}}\exp\Big\{-\frac{2m\omega}{\hbar}r^2\Big\}.
\end{aligned} \tag{42}$$

This is our benchmark.

It can be shown analytically (using a harmonic oscillator basis) that the Hartree (H) solution (remind that there is no exchange because the two particles are different) leads to $E_{int} = \frac{1}{\sqrt{2}}\hbar\omega$ and $\rho^{(l)}_{int}(r) = \sqrt{\frac{m\omega}{\sqrt{2}\pi\hbar}}\exp(-\frac{m\omega}{\sqrt{2}\hbar}r^2)$. Thus, the H energy is $2/\sqrt{2}$ (≈ 1.4) times more important than the one of the benchmark and the density is 1.7 times more spread. The H solution is much more delocalized than the benchmark because the c.m. correlations are neglected [8].

2. *The internal DFT exact functional.*

Applying the multicomponent internal DFT formalism developped in Ref. [17] (whose equations have a relatively similar form than the "one kind of particles" internal DFT ones recalled in §II A) we can rewrite the internal energy as a functional of $\rho^{(1)}_{int}$ and $\rho^{(2)}_{int}$ (we have choosen $v_{int} = 0$)

$$E_{int}[\rho^{(1)}_{int},\rho^{(2)}_{int}] = (\varphi^{(1)}_{int}|\frac{p^2}{2m}|\varphi^{(1)}_{int}) + (\varphi^{(2)}_{int}|\frac{p^2}{2m}|\varphi^{(2)}_{int}) + E^{(12)}_{H}[\rho^{(1)}_{int},\rho^{(2)}_{int}] + E^{(12)}_{potC}[\rho^{(1)}_{int},\rho^{(2)}_{int}] + E^{(12)}_{\Delta kin}[\rho^{(1)}_{int},\rho^{(2)}_{int}], \tag{43}$$

where

$$\begin{aligned}
E^{(12)}_{H}[\rho^{(1)}_{int},\rho^{(2)}_{int}] &= \int drdr'\rho^{(1)}_{int}(r)\rho^{(2)}_{int}(r')\frac{1}{4}m\omega^2(r-r')^2 \\
E^{(12)}_{potC}[\rho^{(1)}_{int},\rho^{(2)}_{int}] &= \int dr\, dr'\gamma^{(12)}_{int}(r,r')\frac{1}{4}m\omega^2(r-r')^2 - E^{(12)}_{H}[\rho^{(1)}_{int},\rho^{(2)}_{int}] \\
E^{(12)}_{\Delta kin}[\rho^{(1)}_{int},\rho^{(2)}_{int}] &= (\psi_{int}|\frac{\tau^2}{2\mu}|\psi_{int}) - (\varphi^{(1)}_{int}|\frac{p^2}{2m}|\varphi^{(1)}_{int}) - (\varphi^{(2)}_{int}|\frac{p^2}{2m}|\varphi^{(2)}_{int}).
\end{aligned} \tag{44}$$

$E^{(12)}_{H}$ is the H energy, $E^{(12)}_{potC}$ is the "standard" correlations energy linked to the particle-particle interaction and $E^{(12)}_{\Delta kin}$ is the energy associated to the correlations contained in the interacting kinetic energy; it is the only term that contains explicitely the c.m. correlations. Using the results of §IV A 1, we can reformulate those functionals as explicit functionals of $\rho^{(1)}_{int}$ and $\rho^{(2)}_{int}$. We have [17]

$$\gamma^{(12)}_{int}(r,r') = \int dr^{(1)}dr^{(2)}\delta(R)|\psi_{int}(r^{(1)}-r^{(2)})|^2\delta\big(r-(r^{(1)}-R)\big)\delta\big(r'-(r^{(2)}-R)\big)$$

---

[8] The c.m. correlations tend to localize the densities compared to the independent particle approximation, which can be understood as follows: if c.m. correlations are taken into account, when one particle moves in a direction where the potential well is higher, the other one will have to move in the opposite direction where the potential well is also higher; thus, the first particle will feel the repulsion present in the independent particle approximation, but also the repulsion that feels the second particle through the c.m. correlations, which will localize its density.

$$= 2|\psi_{int}(2r)|^2\delta(r+r') = \rho^{(l)}_{int}(r)\delta(r+r') = \frac{1}{2}\Big(\rho^{(1)}_{int}(r)+\rho^{(2)}_{int}(r)\Big)\delta(r+r'). \tag{45}$$

Thus (we pose $\varphi^{(l)}_{int}=\sqrt{\rho^{(l)}_{int}}$) [9]

$$E^{(12)}_{potC}[\rho^{(1)}_{int},\rho^{(2)}_{int}] = \int dr\frac{1}{2}\Big(\rho^{(1)}_{int}(r)+\rho^{(2)}_{int}(r)\Big)m\omega^2r^2 - E^{(12)}_{H}[\rho^{(1)}_{int},\rho^{(2)}_{int}] \tag{46}$$

$$E^{(12)}_{\Delta kin}[\rho^{(1)}_{int},\rho^{(2)}_{int}] = -\frac{3}{2}\hbar\omega + \frac{3}{2}\int dr\Big(\rho^{(1)}_{int}(r)+\rho^{(2)}_{int}(r)\Big)m\omega^2r^2. \tag{47}$$

3. The c.m. correlations functional.

We now apply the idea presented in §III with $\Gamma^{aux}(R)=\Big(\frac{K}{\pi}\Big)^{1/4}\exp\Big\{-\frac{K}{2}R^2\Big\}$. We obtain

$$\begin{aligned} E^{(12)}_{cm} = &-\frac{\hbar^2}{2m}\int dr\Big[\varphi^{(1)*}_{int}(r)\Delta_r\varphi^{(1)}_{int}(r)\Big(2\sqrt{\frac{\pi}{K}}|\varphi^{(2)}_{int}(-r)|^2-1\Big)+\varphi^{(2)*}_{int}(r)\Delta_r\varphi^{(2)}_{int}(r)\Big(2\sqrt{\frac{\pi}{K}}|\varphi^{(1)}_{int}(-r)|^2-1\Big)\Big] \\ &-\frac{\hbar^2}{2m}\sqrt{K\pi}\int dr|\varphi^{(1)}_{int}(r)|^2|\varphi^{(2)}_{int}(-r)|^2, \end{aligned} \tag{48}$$

and for the local c.m. correlations potentials ($l=1,2$)

$$\begin{aligned} U^{(l)}_{cm}(r) = &-\frac{1}{\varphi^{(l)}_{int}(r)}\frac{\hbar^2}{2m}\Big(2\sqrt{\frac{\pi}{K}}|\varphi^{(m\neq l)}_{int}(-r)|^2-1\Big)\Delta_r\varphi^{(l)}_{int}(r) \\ &-\frac{\hbar^2}{2m}\Big(2\sqrt{\frac{\pi}{K}}\varphi^{(m\neq l)*}_{int}(-r)\Delta_r\varphi^{(m\neq l)*}_{int}(-r)+\sqrt{K\pi}|\varphi^{(m\neq l)}_{int}(-r)|^2\Big). \end{aligned} \tag{49}$$

4. Numerical results.

We use a unit system where $\hbar=m=1$ and we choose $\omega=1$. Table I and Fig. 1 give numerical results for the following formalisms:

- Benchmark (described in §IV A 1),
- $E^{(12)}_{H}$, called "H only",
- $E^{(12)}_{H}+E^{(12)}_{C}$, called "H + standard correlations",
- $E^{(12)}_{H}+E^{(12)}_{C}-<\frac{\mathbf{P}^2}{2mN}>$, called "H + standard correlations + standard c.m. correction",
- $E^{(12)}_{H}+E^{(12)}_{C}+E^{(12)}_{\Delta kin}$, called "exact internal DFT",
- $E^{(12)}_{H}+E^{(12)}_{C}+E_{cm}$, called "internal DFT with c.m. correlations functional"; we obtain $K\approx 3.9$ with the method described in §III D.

First of all, we see from table I and Fig. 1 that "exact internal DFT" reproduces perfectly the total energy and the densities of the benchmark, so that the non-interacting v-representability [15, 21] is perfectly achieved. This is not a surprise because the non-interacting v-representability is trivially achieved when one deals with only one particle of each kind (because it is always possible to reach $\varphi^{(l)}_{int}=\sqrt{\rho^{(l)}_{int}}$).

---

[9] Note that, even if in the general case the functional $E^{(12)}_{potC}+E^{(12)}_{\Delta kin}$ is universal [15, 16], the forms (46) and (47) are limited to the two different particles case because exchange effects are not taken into account; they thus cannot be used to describe a system composed by and arbitrary number of particles of each kind. The universal functional, applicable to an arbitrary number of particles, is more involved but should permit to recover (46) and (47) in the limit of a system composed by two different particles.

Formalism	Non-interacting kin. energy	$E_H^{(12)}$	$E_C^{(12)}$	$- < \frac{\mathbf{P}^2}{2mN} >$ or $E_{\Delta kin}^{(12)}$ or E_{cm}	Total energy
H only	0.353	0.353	0	0	0.71
H + stand. corr.	0.5	0.25	0.25	0	1.00
H + stand. corr. + stand. c.m. correct.	0.706	0.177	0.177	-0.353	0.71
Exact internal DFT	1.000	0.125	0.125	-0.750	0.50
Internal DFT with c.m. corr. ft	1.225	0.120	0.120	-0.918	0.55

TABLE I: The energies of the various formalisms (in units where $\hbar = m = 1$; benchmark: total energy $= 0.50$ and interacting kinetic energy $(\psi_{int}|\frac{\tau^2}{2\mu}|\psi_{int}) = 0.25$).



FIG. 1: (Coloured version online.) The internal densities $\rho_{int}^{(l)}$ of the various formalisms (x-axis: position in units where $\hbar = m = 1$).

From table I, we see that the non-interacting kinetic energies of the various mean-field like calculations *cannot* be compared to the interacting kinetic energy In particular, there is a factor 4 between the "exact internal DFT" non-interacting kinetic energy (equal to 1) and the interacting kinetic energy (equal to 0.250). Indeed, as already discussed in §II A, it is the "non-interacting kinetic energy + $E_{\Delta kin}$" that is comparable to the interacting kinetic energy. Exact internal DFT then reaches perfectly the benchmark: $1.000 - 0.750 = 0.250$. Internal DFT with E_{cm} gives $1.225 - 0.928 = 0.307$, which reproduces fairly well the benchmark considering that the "standard" correlations part of $E_{\Delta kin}^{(12)}$ has been neglected. The result with standard c.m. correction gives $0.706 - 0.353 = 0.353$, which is worse.

From the point of view of the total energy, "H only" is far away from the benchmark, as well as "H + stand. corr." and "H + stand. corr. + stand. c.m. correct.". "Internal DFT with c.m. corr. ft." is the closest to the benchmark.

From the point of view of the densities, Fig. 1 shows that "internal DFT with c.m. corr. ft." is very close from the benchmark and represents a great improvement compared to the other results. Nevertheless, we see some differences that should mostly be due to the fact that the "standard" correlations part of $E_{\Delta kin}^{(12)}$ is not taken into account. In realistic mean-field like calculations, however, this "problem" should not appear because those "standard" correlations are mostly taken into account in the parametrized functionals that are commonly used.

B. Model system of identical Bosons (smooth interaction).

We now study the results of internal DFT on another 1D system, with a "smooth" interaction. We consider a system composed of N identical Bosons of mass m and positions $\{r_i\}$, without spin, in a condensate state and with an attractive 2-body interaction of the form ($e > 0$)

$$u(r-r') = -\frac{1}{\sqrt{(r-r')^2+e}}. \tag{50}$$

The more e is large, the more the potential is smooth. This allows to model, in some way, some features of ^{4}He droplets.

The internal DFT energy functional is given by (φ_{int} is the 1-body orbital describing the bosons with $\rho_{int} = N|\varphi_{int}|^2$ and we consider $v_{int} = 0$)

$$E_{int}[\rho_{int}] \ = \ N(\varphi_{int}|\frac{p^2}{2m}|\varphi_{int}) + E_H[\rho_{int}] \times (1-\frac{1}{N}) + E_C^{stand}[\rho_{int}] + E_{cm}[\rho_{int}], \tag{51}$$

where $(1-\frac{1}{N})E_H[\rho_{int}]$ represents the H energy where the self-interaction has been substracted and $E_C^{stand}[\rho_{int}]$ is the "standard" correlations energy defined as

$$E_C^{stand}[\rho_{int}] = \frac{1}{2}\int dr\, dr'\, \gamma_{int}(r,r')\, u(r-r') - E_H[\rho_{int}] \times (1-\frac{1}{N}) + E_{\Delta kin}[\rho_{int}] - E_{cm}[\rho_{int}]. \tag{52}$$

The c.m. correlations energy is defined as in §III F, where Γ^{aux} is defined as in §III D

$$\begin{aligned} E_{cm}[\rho_{int}] \ &= \ -\frac{\hbar^2}{2m}\int dr\ \sqrt{\rho_{int}(\mathbf{r})}\Delta_{\mathbf{r}}\sqrt{\rho_{int}(\mathbf{r})} \times \Big(\sqrt{\frac{\pi}{K(N)}}\int dr'\ \frac{1}{N}\rho_{int}(\mathbf{r}')f_2[\rho_{int}](r+r') - 1\Big) \\ &\quad -\frac{\hbar^2}{2mN}\sqrt{\pi K(N)}\int dr\ |\varphi_{int}(r)|^2 \int dr'\ \frac{1}{N}\rho_{int}(\mathbf{r}') \times f_2[\rho_{int}](r+r'), \end{aligned}$$

and f_2 is defined by Eq. (37). The internal KS equation is

$$\Big(-\frac{\hbar^2}{2m}\Delta + U_H[\rho_{int}] \times (1-\frac{1}{N}) + U_C^{stand}[\rho_{int}] + U_{cm}[\rho_{int}]\Big)\varphi_{int} = \epsilon\varphi_{int}, \tag{53}$$

where $U_C^{stand} = \delta E_C^{stand}/\delta\rho_{int}$ and

$$\begin{aligned} U_{cm}[\rho_{int}](r) \ &= \ -\frac{\hbar^2}{2m}\Big\{\frac{1}{\sqrt{\rho_{int}(r)}}\Delta_r\sqrt{\rho_{int}(r)} \times \Big(\sqrt{\frac{\pi}{K(N)}}\int dr'\frac{1}{N}\rho_{int}(\mathbf{r}')f_2[\rho_{int}](r+r') - 1\Big) \\ &\quad + \sqrt{\frac{\pi}{K(N)}}\ \int dr'\ \sqrt{\rho_{int}(\mathbf{r}')}\Delta_{\mathbf{r}}\sqrt{\rho_{int}(\mathbf{r}')} \times \frac{N-1}{N}f_2[\rho_{int}](r+r')\Big\} \\ &\quad -\frac{\hbar^2}{2mN}\sqrt{K(N)\pi}\int dr'\frac{1}{N}\rho_{int}(\mathbf{r}') \times f_2[\rho_{int}](r+r'). \end{aligned} \tag{54}$$

Formalism	Non-interacting kin. energy	$E_H \times (1-\frac{1}{N})$	$- < \frac{\mathbf{P}^2}{2mN} >$ or E_{cm}	Total energy
H only	0.133	-0.626	0	-0.49
H + stand. c.m. correct.	0.260	-0.712	-0.065	-0.52
Internal DFT with c.m. corr. ft	0.535	-0.776	-0.418	-0.66

TABLE II: The energies of the various formalisms in the $N = 2$ case (in units where $\hbar = m = 1$; benchmark: total energy $= -0.59$ and interacting kinetic energy $(\psi_{int}|\frac{\tau^2}{2\mu}|\psi_{int}) = 0.12$).

For the $N = 2$ case, we can compute a benchmark. Indeed, by use of Jacobi coordinates, the internal Hamiltonian can be rewritten $H_{int} = \frac{\tau^2}{2\mu} - \frac{1}{\sqrt{\xi^2+e}}$, where $\mu = m/2$ is the reduced mass. It is then possible to calculate numerically the exact many body solution ground state ψ_{int} and the c.m. frame one-body density $\rho_{int}(r) = 4|\psi_{int}(2r)|^2$.

In the following, we neglect E_C^{stand} because we have no simple way to evaluate it as a functional of ρ_{int} and it should be a correction in the general case (even if not always fully negligible). The next results are given for:



FIG. 2: (Coloured version online.) The internal density $\rho_{int}/2$ of the various formalisms in the $N = 2$ bosonic case (x-axis: position in units where $\hbar = m = 1$).

$N =$	Non-interacting kin. energy	Total energy
2	0.133	-0.491
3	0.385	-1.75
4	0.717	-3.79
5	1.138	-6.66
6	1.632	-10.37

TABLE III: The "H only" energies for various N (in units where $\hbar = m = 1$).

- $E_H \times (1 - \frac{1}{N})$, called "$H$ only",
- $E_H \times (1 - \frac{1}{N}) - < \frac{\mathbf{P}^2}{2mN} >$, called "H + standard c.m. correction",
- $E_H \times (1 - \frac{1}{N}) + E_{cm}$, called "internal DFT with c.m. correlations functional".

Table II and Fig. 2 show the energies and densities in the $N = 2$ case. We see that "internal DFT with c.m. corr. ft." reproduces fairly well the benchmark, at least much better than "H only" and "H + stand. c.m. correct" (we

$N =$	Non-interacting kin. energy	E_{cm}	Total energy	K	Interacting kin. energy
2	0.535	-0.418	-0.66	1.94	0.117
3	0.463	-0.185	-1.90	1.74	0.278
4	0.702	-0.196	-3.97	2.74	0.507
5	1.014	-0.217	-6.84	4.01	0.799
6	1.390	-0.239	-10.56	4.54	1.151

TABLE IV: The "internal DFT with c.m. corr. ft." energies for various N (in units where $\hbar = m = 1$).



FIG. 3: (Coloured version online.) The internal density $\rho_{int}/6$ of the various formalisms in the $N = 6$ bosonic case (x-axis: position in units where $\hbar = m = 1$).



FIG. 4: (Coloured version online.) f_2 for $N = 3$ to 6 ($N = 2$ is a delta function; x-axis: position in units where $\hbar = m = 1$.

nevertheless see some differences, which should mostly be due to the fact that E_C^{stand} has been neglected). Again, we see that the internal DFT non-interacting kinetic energy *cannot* be compared to the interacting kinetic energy. It is the internal DFT "non-interacting kinetic energy + E_{cm}" ($0.535 - 0.418 = 0.117$) that is comparable to the interacting kinetic energy (0.12), as underlined in §II A. The result with standard c.m. correction gives $0.260 - 0.065 = 0.195$, which is worse.

Tables III and IV show that "H only" total energy becomes closer to internal DFT total energy as N grows. Moreover, table IV shows that the part of the c.m. correlations energy in the total internal DFT energy decreases as N grows (63% for $N = 2$; 2% for $N = 6$). As a consequence, and even is shell effets play a role, the internal DFT non-interacting kinetic energy tends to become closer to the interacting kinetic energy as N grows (factor 4.6 for $N = 2$; factor 1.2 for $N = 6$). The "H only" and internal DFT densities also becomes closer as N grows, see Fig. 3 for $N = 5$, as awaited.

Finally, we see from table IV that K grows as N grows and from Fig. 4 that the maximum value of f_2 also grows as N grows (but the maximum value $\frac{1}{N^D} f_2$ diminishes), confirming the reasonings of §III D and Appendix C.

V. CONCLUSION.

Internal DFT provides an existence theorem for a c.m. correlations energy functional associated to a local potential.

In this article, we have constructed such a functional, without including any free parameter. The use of this functional is justified by a strong formal background and variants suitable for fermionic as well as bosonic systems have been proposed. The resulting scheme is numerically manageable and represents a well founded alternative to projection techniques to treat the c.m. correlations. It can directly be added to actual energy functionals (but a reffiting of them then should be necessary). Moreover, it permits to recover clearly and precisely the value of the interacting kinetic energy and represents a manageable way to include the c.m. correlations in time-dependent calculations of self-bound systems.

We have presented first convincing numerical results on 1D model systems. They amongst other show that the developped functional represents a great improvement compared to the "standard c.m. correction" commonly used in nuclear Physics (of the form $- < \frac{\mathbf{P}^2}{2mN} >$), especially from the point of view of the energies. The next step will be to include the proposed c.m. correlations functional in realistic 3D calculations, for instance in mean-field like calculations of nuclei with Skyrme interaction [2, 9]. As the "standard" correlations are mostly taken into account in the commonly used functionals, the 3D results should be even more convincing than the 1D ones.

Finally, even if the proposed Gaussian form for $\Gamma^{aux}(\mathbf{R})$ has been proved to be pertinent, it could be interesting to search for other forms, i.e. with other variations around $\mathbf{R} = \mathbf{0}$, to obtain the most precise description of atomic nuclei, Helium droplets or molecular systems with explicit treatment of the nuclei.

Acknowledgments.

The author is particularly grateful to T. Duguet for numerous enlightening discussions and reading of the manuscript. The author also gratefully acknowledges the Conseil Régional d'Aquitaine for support.

Appendix A: Details of the calculation that leads to E_{cm}.

We evaluate $\int d\mathbf{r}_1 \ldots d\mathbf{r}_N \delta(\mathbf{R}) \psi^*_{int}(\mathbf{r}_1, \ldots, \mathbf{r}_N) \sum_{i=1}^{N} \frac{\mathbf{p}_i^2}{2m} \psi_{int}(\mathbf{r}_1, \ldots, \mathbf{r}_N)$ using the approximation (17) for ψ_{int} and obtain [10]

$$
\begin{aligned}
\int \quad & d\mathbf{r}_1 \ldots d\mathbf{r}_N \delta(\mathbf{R}) \frac{1}{\Gamma^{aux*}(\mathbf{R})} \psi^{aux*}(\mathbf{r}_1, \ldots, \mathbf{r}_N) \sum_{i=1}^{N} \frac{\mathbf{p}_i^2}{2m} \frac{1}{\Gamma^{aux}(\mathbf{R})} \psi^{aux}(\mathbf{r}_1, \ldots, \mathbf{r}_N) \qquad \text{(A1)} \\
= & -\frac{\hbar^2}{2m} \frac{1}{|\Gamma^{aux}(\mathbf{0})|^2} \frac{1}{N!} \sum_{P,P'} (-1)^{p+p'} \sum_{i=1}^{N} \int d\mathbf{r}_i \; F_i^{P,P'}[\{\varphi^k_{int}\}](\mathbf{r}_i) \times \varphi^{P(i)*}_{int}(\mathbf{r}_i) \Delta_{\mathbf{r}_i} \varphi^{P'(i)}_{int}(\mathbf{r}_i) \\
& -\frac{\hbar^2}{2m} \frac{1}{N^2} \frac{1}{\Gamma^{aux*}(\mathbf{0})} \Delta_{\mathbf{R}} \frac{1}{\Gamma^{aux}(\mathbf{R})} \Big|_{\mathbf{R}=\mathbf{0}} \frac{1}{N!} \sum_{P,P'} (-1)^{p+p'} \sum_{i=1}^{N} \int d\mathbf{r}_i \; F_i^{P,P'}[\{\varphi^k_{int}\}](\mathbf{r}_i) \times \varphi^{P(i)*}_{int}(\mathbf{r}_i) \varphi^{P'(i)}_{int}(\mathbf{r}_i) \\
= & -\frac{\hbar^2}{2m} \frac{1}{|\Gamma^{aux}(\mathbf{0})|^2} \sum_{i=1}^{N} \Big\{ \int d\mathbf{r} \; F_i[\{\varphi^{k \neq i}_{int}\}](\mathbf{r}) \times \varphi^{i*}_{int}(\mathbf{r}) \Delta_{\mathbf{r}} \varphi^{i}_{int}(\mathbf{r}) \\
& \qquad + \frac{1}{N!} \sum_{P,P' \neq P} (-1)^{p+p'} \int d\mathbf{r} \; F_i^{P,P' \neq P}[\{\varphi^k_{int}\}](\mathbf{r}) \times \varphi^{P(i)*}_{int}(\mathbf{r}) \Delta_{\mathbf{r}} \varphi^{P'(i)}_{int}(\mathbf{r}) \Big\} \\
& -\frac{\hbar^2}{2mN} \frac{1}{\Gamma^{aux*}(\mathbf{0})} \Delta_{\mathbf{R}} \frac{1}{\Gamma^{aux}(\mathbf{R})} \Big|_{\mathbf{R}=\mathbf{0}} \frac{1}{N} \sum_{i=1}^{N} \Big\{ \int d\mathbf{r} \; F_i[\{\varphi^{k \neq i}_{int}\}](\mathbf{r}) \times |\varphi^{i}_{int}(\mathbf{r})|^2 \\
& \qquad + \frac{1}{N!} \sum_{P,P' \neq P} (-1)^{p+p'} \int d\mathbf{r} \; F_i^{P,P' \neq P}[\{\varphi^k_{int}\}](\mathbf{r}) \times \varphi^{P(i)*}_{int}(\mathbf{r}) \varphi^{P'(i)}_{int}(\mathbf{r}) \Big\},
\end{aligned}
$$

---

[10] Note that $\Delta_{\mathbf{r}_i} \frac{1}{\Gamma^{aux}(\mathbf{R})} = \frac{1}{N^2} \Delta_{\mathbf{R}} \frac{1}{\Gamma^{aux}(\mathbf{R})}$, which is obtained by a simple variable change and is usefull for the calculation.

where we have defined ($D = 1$, 2 or 3 is the dimension in which the calculation is done)

$$F_i^{P,P'}[\{\varphi^k_{int}\}](\mathbf{r}) \;=\; N^D \int \Pi_{\substack{j=1\\ j\neq i}}^N d\mathbf{r}_j \delta\Big(\sum_{\substack{k=1\\ k\neq i}}^N \mathbf{r}_k + \mathbf{r}\Big) \Pi_{\substack{j=1\\ j\neq i}}^N \varphi_{int}^{P(j)*}(\mathbf{r}_j)\varphi_{int}^{P'(j)}(\mathbf{r}_j), \tag{A2}$$

and its diagonal part

$$F_i[\{\varphi^{k\neq i}_{int}\}](\mathbf{r}) = \frac{1}{N!}\sum_P F_i^{P,P}[\{\varphi^k_{int}\}](\mathbf{r}) = N^D \int \Pi_{\substack{j=1\\ j\neq i}}^N d\mathbf{r}_j \delta\Big(\sum_{\substack{k=1\\ k\neq i}}^N \mathbf{r}_k + \mathbf{r}\Big) \Pi_{\substack{j=1\\ j\neq i}}^N |\varphi^j_{int}(\mathbf{r}_j)|^2. \tag{A3}$$

F_i has the following properties: it is real and has the dimension of a density; it is normalized to N^D, i.e. $\int d\mathbf{r} F_i(\mathbf{r}) = N^D$; $\lim_{\mathbf{r}\to\pm\infty} F_i(\mathbf{r}) = 0$; it is a "multiconvolution" of all single densities, unless the one associated to orbital i. The first three points permit to explicit the physical meaning of $\frac{1}{N^D}F_i(\mathbf{r})$: it is the probability that particle i has position $\mathbf{r}$, according to the c.m. coupling with every other particles and their probability distributions.

$F_{P,P'\neq P}$ is only due to exchange effects. In the following, as explained in §II B and §III B, we neglect the pure exchange effects and thus $F_{P,P'\neq P}$. We obtain

$$\begin{aligned}\int\quad & d\mathbf{r}_1 \ldots d\mathbf{r}_N \delta(\mathbf{R})\psi^*_{int}(\mathbf{r}_1,\ldots,\mathbf{r}_N)\sum_{i=1}^N \frac{p_i^2}{2m}\psi_{int}(\mathbf{r}_1,\ldots,\mathbf{r}_N) \\ & \to -\frac{\hbar^2}{2m}\frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}\sum_{i=1}^N \int d\mathbf{r}\; F_i[\{\varphi^{k\neq i}_{int}\}](\mathbf{r}) \times \varphi^{i*}_{int}(\mathbf{r})\Delta_{\mathbf{r}}\varphi^i_{int}(\mathbf{r}) \\ & \quad -\frac{\hbar^2}{2mN}\frac{1}{\Gamma^{aux*}(\mathbf{0})}\Delta_{\mathbf{R}}\frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}}\frac{1}{N}\sum_{i=1}^N \int d\mathbf{r}\; F_i[\{\varphi^{k\neq i}_{int}\}](\mathbf{r}) \times |\varphi^i_{int}(\mathbf{r})|^2.\end{aligned} \tag{A4}$$

We now insert this result in $E_{\Delta kin}$, Eq. (10), and keep only the real part, i.e. $\Re e(E_{\Delta kin})$, as justified in §III B. It only remains the c.m. correlations contribution:

$$\begin{aligned} E_{\Delta kin} \to E_{cm} \;=\; & -\frac{\hbar^2}{2m}\sum_{i=1}^N \int d\mathbf{r}\; \Big(\frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}F_i[\{\varphi^{k\neq i}_{int}\}](\mathbf{r}) - 1\Big) \times \varphi^{i*}_{int}(\mathbf{r})\Delta_{\mathbf{r}}\varphi^i_{int}(\mathbf{r}) \\ & -\frac{\hbar^2}{2mN}\Re e\Big(\frac{1}{\Gamma^{aux*}(\mathbf{0})}\Delta_{\mathbf{R}}\frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}}\Big)\int d\mathbf{r}\; F_i[\{\varphi^{k\neq i}_{int}\}](\mathbf{r}) \times |\varphi^i_{int}(\mathbf{r})|^2 \\ & + PureImaginary[\{\varphi^k_{int}\}], \end{aligned} \tag{A5}$$

where

$$PureImaginary[\{\varphi^k_{int}\}] = i\frac{\hbar^2}{2m}\Im m\sum_{i=1}^N \int d\mathbf{r}\; \frac{1}{|\Gamma^{aux}(\mathbf{0})|^2}F_i[\{\varphi^{k\neq i}_{int}\}](\mathbf{r}) \times \varphi^{i*}_{int}(\mathbf{r})\Delta_{\mathbf{r}}\varphi^i_{int}(\mathbf{r}) \tag{A6}$$

is a pure imaginary functional which counter acts the pure imaginary part of the first line of (A5).

F_i is interesting to underline the Physics in energy considerations. But it is not a fundamental quantity for the potential (that will be obtained by variation of E_{cm}). We thus introduce a more fundamental quantity, which will appear in both the c.m. correlations energy and potential, namely the "two particles c.m. correlations functional" defined in Eq. (20), which is linked to F_i by the relations

$$\begin{aligned} \forall l \neq i: \qquad & F_i[\{\varphi^{k\neq i}_{int}\}](\mathbf{r}') = \int d\mathbf{r}|\varphi^l_{int}(\mathbf{r})|^2 \times f_{i,l\neq i}[\{\varphi^{k\neq i,l}_{int}\}](\mathbf{r}+\mathbf{r}'), \\ & \frac{\delta F_i[\{\varphi^{k\neq i}_{int}\}](\mathbf{r}')}{\delta\varphi^{l*}_{int}(\mathbf{r})} = \varphi^l_{int}(\mathbf{r}) \times f_{i,l\neq i}[\{\varphi^{k\neq i,l}_{int}\}](\mathbf{r}+\mathbf{r}'). \end{aligned} \tag{A7}$$

When (A7) is inserted in (A5), we obtain the form (18) for the c.m. correlation energy, presented in §III B.

Appendix B: The multiconvolution theorem.

We define the Fourier transform $\mathcal{T}$ of an integrable function $L : \Re e \to \Im m$ as

$$\forall \mathbf{r}, \mathbf{s} \in \Re e : \quad \mathcal{T}[L](\mathbf{s}) = \int d\mathbf{r}\; e^{-2\pi i \mathbf{s}.\mathbf{r}}\; L(\mathbf{r}) \tag{B1}$$

and the inverse Fourier transform $\mathcal{T}^{-1}$ of a function $\tilde{L} : \Re e \to \Im m$ as

$$\forall \mathbf{r}, \mathbf{s} \in \Re e : \quad \mathcal{T}^{-1}[\tilde{L}](\mathbf{r}) = \int d\mathbf{s}\; e^{2\pi i \mathbf{s}.\mathbf{r}}\; \tilde{L}(\mathbf{s}).$$

We start from $(K+1)$ integrable functions $g_i : \Re e \to \Im m$ and define the "multiconvolution"

$$C[\{g_i\}](\tilde{\mathbf{r}}) = \int d\mathbf{r}_1 \dots d\mathbf{r}_K\; g_1(\mathbf{r}_1) \times \dots \times g_K(\mathbf{r}_K) \times g_{K+1}(-\sum_{i=1}^{K} \mathbf{r}_i - \tilde{\mathbf{r}}).$$

We calculate the Fourier transform of C:

$$\begin{aligned}
\mathcal{T}[C](\mathbf{s}) &= \int d\tilde{\mathbf{r}}\; e^{-2\pi i \mathbf{s}.\tilde{\mathbf{r}}}\; C[\{g_i\}](\tilde{\mathbf{r}}) \\
&= \int d\tilde{\mathbf{r}}\; d\mathbf{r}_1 \dots d\mathbf{r}_K\; e^{-2\pi i \mathbf{s}.\tilde{\mathbf{r}}}\; g_1(\mathbf{r}_1) \times \dots \times g_K(\mathbf{r}_K) \times g_{K+1}(-\sum_{i=1}^{K} \mathbf{r}_i - \tilde{\mathbf{r}}) \\
&= \int d\mathbf{w}\; d\mathbf{r}_1 \dots d\mathbf{r}_K\; e^{-2\pi i \mathbf{s}.(-\sum_{i=1}^{K} \mathbf{r}_i - \mathbf{w})}\; g_1(\mathbf{r}_1) \times \dots \times g_K(\mathbf{r}_K) \times g_{K+1}(\mathbf{w}) \\
&= \Pi_{i=1}^{K+1} \int d\mathbf{r}_i\; e^{2\pi i \mathbf{s}.\mathbf{r}_i} g_i(\mathbf{r}_i) \\
&= \Pi_{i=1}^{K+1} \mathcal{T}[g_i](-\mathbf{s}),
\end{aligned} \tag{B2}$$

where the second line has been obtained doing the variable change $\mathbf{w} = -\sum_{i=1}^{K} \mathbf{r}_i - \tilde{\mathbf{r}}$, which implies $d\mathbf{w} = -d\tilde{\mathbf{r}}$.

This is the "convolution theorem" [36] generalized to multiconvolutions, which states that the Fourier transform of a multiconvolution is the product of the Fourier transforms of each functions that enter into the multiconvolution. This theorem permits to fasten the numerical calculation of $C[\{g_i\}](\tilde{\mathbf{r}})$. Indeed, it translates a multiconvolution to a multiplication of functions. Once every $\mathcal{T}[g_i]$ are calculated, $C[\{g_i\}]$ is given by the inverse Fourier transform of their direct product:

$$C[\{g_i\}](\tilde{\mathbf{r}}) = \mathcal{T}^{-1}\Big[\Pi_{i=1}^{K+1} \mathcal{T}[g_i](-\mathbf{s})\Big](\tilde{\mathbf{r}}) = \mathcal{T}^{-1}\Big[\Pi_{i=1}^{K+1} \mathcal{T}[g_i]\Big](-\tilde{\mathbf{r}}).$$

Note that this relationship is only valid for the form (B1) of the Fourier transform. For forms normalized in other ways, a constant scaling factor, typically proportional to $\Big[(2\pi)^{D/2}\Big]^K$, will appear.

Appendix C: Some properties of Γ^{aux} when N becomes very large.

The limit where the c.m. correlations become negligible is obtained when N becomes very large, as mentioned in §III C. Indeed, the "two particles c.m. correlations functional" $f_{i,l\neq i}$ then tends to become constant and delocalized in the whole space. We note

$$\lim_{N \to +\infty} f_{i,l\neq i} = Constant. \tag{C1}$$

The normalization condition (24) thus implies, when N is very large (remind that Γ^{aux} is implicitely dependent of N)

$$\lim_{N \to +\infty} |\Gamma^{aux}(\mathbf{0})|^2 = \lim_{N \to +\infty} \int d\mathbf{r}\; d\mathbf{r}'\; |\varphi^i_{int}(\mathbf{r})|^2 |\varphi^{l\neq i}_{int}(\mathbf{r}')|^2 f_{i,l\neq i}[\{\varphi^{k\neq i,l}_{int}\}](\mathbf{r}+\mathbf{r}') = Constant. \tag{C2}$$

When those results are inserted in E_{cm}, Eq. (18), we see that its first and obviously third lines become null, and that its second line becomes proportional to $\frac{1}{N} \times \frac{1}{\Gamma^{aux*}(\mathbf{0})} \Delta_{\mathbf{R}} \frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}} \times |\Gamma^{aux}(\mathbf{0})|^2$, which must tend to zero so that E_{cm} can be neglected, as it should be when N becomes very large. This implies

$$\lim_{N \to +\infty} \frac{1}{N} \times \frac{1}{\Gamma^{aux*}(\mathbf{0})} \Delta_{\mathbf{R}} \frac{1}{\Gamma^{aux}(\mathbf{R})}\Big|_{\mathbf{R}=\mathbf{0}} \times |\Gamma^{aux}(\mathbf{0})|^2 \to 0. \tag{C3}$$

To obtain a second relation, we note $\mathcal{R}$ the region of space where the system has a non zero density and V the corresponding volume. For very large systems, we have

$$\begin{aligned} |\varphi^i_{int}(\mathbf{r})|^2 &\approx \tfrac{1}{V}, \qquad \text{for } \mathbf{r} \in \mathcal{R}, \\ &\approx 0, \qquad \text{for } \mathbf{r} \notin \mathcal{R}. \end{aligned} \tag{C4}$$

Inserting those results in the definition (23) of $f_{i,l\neq i}$ gives

$$\begin{aligned} f_{i,l\neq i}(\mathbf{r}) &\approx \tfrac{N^D}{V}, \qquad \text{for } \mathbf{r} \in \mathcal{R}, \\ &\approx 0, \qquad \text{for } \mathbf{r} \notin \mathcal{R}. \end{aligned} \tag{C5}$$

In the general case, we have $V < kN$, where k is a constant (it is for saturating systems, as nuclear ones [2], that V becomes close, but still inferior, to kN). Thus

$$\lim_{N \to +\infty} f_{i,l\neq i}(\tilde{\mathbf{r}}) = +\infty, \tag{C6}$$

whatever the dimension in which the calculation is done (but $\lim_{N \to +\infty} \frac{1}{N^D} f_{i,l\neq i}(\mathbf{r}) \to \lim_{N \to +\infty} \frac{1}{V}$, which tends to zero). As a consequence of Eqs. (C1), (C2) and (C6), we deduce a second relation that Γ^{aux} should satisfy:

$$\lim_{N \to +\infty} |\Gamma^{aux}(\mathbf{0})|^2 \to +\infty. \tag{C7}$$

---

[1] K. W. Schmid, Eur. Phys. J. A **12**, 29 (2001).
[2] P. Ring and P. Schuck, *The Nuclear Many-Body Problem*, Springer Verlag, New York, Heidelberg, Berlin, (1980).
[3] R. E. Peierls and J. Yoccoz, Proc. Phys. Soc. **70**, 381 (1957).
[4] R. E. Peierls and D. J. Thouless, Nucl. Phys. **38**, 154 (1962).
[5] S. Marcos, H. Flocard and P. H. Heenen, Nucl. Phys. A **410**, 125 (1983).
[6] S. Marcos, H. Flocard and P. H. Heenen, Phys. Lett. **134B**, 287 (1983).
[7] K. W. Schmid and P. G. Reinhard, Nucl. Phys. A **530**, 283 (1991).
[8] J. Dobaczewski, J. Phys. G: Nucl. Part. Phys **36**, 105105 (2009).
[9] M. Bender, P.-H. Heenen and P.-G. Reinhard, Rev. Mod. Phys. **75**, 121 (2003).
[10] R. R. Rodríguez-Guzmán and K. W. Schmid, Eur. Phys. J. A **19**, 45 (2004); *ibid.* 61 (2004).
[11] M. Bender, K. Rutz, P.-G. Reinhard and J.A. Maruhn, Eur. Phys. J A **7**, 467 (2000).
[12] A. S. Umar, V. E. Oberacker, J. Phys. G : Nucl. Part. Phys. **36**, 025101 (2009).
[13] J. M. Irvine, J. Phys. G : Nucl. Phys. **6**, L175 (1980).
[14] K. W. Schmid, Eur. Phys. J. A **12**, 29 (2001).
[15] J. Messud, M. Bender and E. Suraud, Phys. Rev. C **80**, 054314 (2009).
[16] J. Messud, Phys. Rev. C **80**, 054614 (2009).
[17] J. Messud, Phys. Rev. A **84**, 052113 (2011).
[18] P. Hohenberg and W. Kohn, Phys. Rev. **136**, 864 (1964).
[19] W. Kohn and L. J. Sham, Phys. Rev. **140**, 1133 (1965).
[20] W. Kohn, Rev. Mod. Phys. **71**, 1253 (1999).
[21] R. M. Dreizler and E. K. U. Gross, *Density Functional Theory*, (Springer, Berlin, 1990)
[22] B. G. Giraud, Phys. Rev. C **77**, 014311 (2008).
[23] K. Yabana, Prog. Theor. Phys. **76**, 414 (1986).
[24] K. Dietrich, Nucl. Phys. **A606**, 63 (1996).
[25] J. Engel, Phys. Rev. C **75**, 014306 (2007).
[26] N. Barnea, Phys. Rev. C **76**, 067302 (2007).
[27] M. Barranco, R. Guardiola, S. Hernańdez, R. Mayol, J. Navarro and M. Pi, J. Low Temp. Phys. **142**, 1 (2006).

[28] R. T. Sharp, G. K. Horton, Phys. Rev. **90**, 317 (1953).
[29] J. D. Talman, W. F. Shadwick, Phys. Rev. A **14**, 36 (1976).
[30] S. Kümmel, L. Kronik, Rev. Mod. Phys. **80**, 3 (2008).
[31] J.B. Krieger, Y. Li, G. J. Iafrate, Phys. Rev. A **45** (1992) 101.
[32] J.B. Krieger, Y. Li, G. J. Iafrate, Phys. Rev. A **46** (1992) 5453.
[33] J. Messud, P. M. Dinh, P.-G. Reinhard and E. Suraud, Ann. Phys. (Berlin) **523** (2011) 270
[34] C. A. Ullrich, U. J. Gossmann, E. K. U. Gross, Phys. Rev. Lett. **74** (1995) 872.
[35] R. G. Parr and W. Yang, *Density-Functional Theory of Atoms and Molecules*, Oxford University Press (1989)
[36] Y. Katznelson, *An Introduction to Harmonic Analysis*, (Cambridge University Press, 2004).